

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>C12Q 1/68</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/61817</b> <b>(43) International Publication Date:</b> 19 October 2000 (19.10.2000)
<b>(21) International Application Number:</b> PCT/US00/09742 <b>(22) International Filing Date:</b> 12 April 2000 (12.04.2000) <b>(30) Priority Data:</b> 09/290,452 12 April 1999 (12.04.1999) US <b>(60) Parent Application or Grant</b> NANOGEN/BECTON DICKINSON PARTNERSHIP [/]; O. EDMAN, Carl, F. [/]; O. NERENBURG, Michael, I. [/]; O. WESTIN, Lorelei, P. [/]; O. CARRINO, John, J. [/]; O. EDMAN, Carl, F. [/]; O. NERENBURG, Michael, I. [/]; O. WESTIN, Lorelei, P. [/]; O. CARRINO, John, J. [/]; O. MURDOCK, Douglas, C. ; O.	<b>Published</b>	
<b>(54) Title: MULTIPLEX AMPLIFICATION AND SEPARATION OF NUCLEIC ACID SEQUENCES ON A BIOELECTRONIC MICROCHIP USING ASYMMETRIC STRUCTURES</b> <b>(54) Titre: AMPLIFICATION MULTIPLEX ET SEPARATION DE SEQUENCES D'ACIDE NUCLEIQUE SUR UNE MICRO-PUCE BIOELECTRONIQUE AU MOYEN DE STRUCTURES ASYMETRIQUES</b>		
<b>(57) Abstract</b> <p>A method for amplifying nucleic acids is provided wherein detection of amplified species is enhanced by the use of asymmetric amplifications. Such amplification is made asymmetric by using divergent ratios of amplification primers or by using non-extending and/or non-cleavable amplification primers. Detection of the amplicons is improved because maintenance of single stranded species of amplicons during amplification facilitates their direct capture by immobilized probes without having to include denaturing steps.</p> <b>(57) Abrégé</b> <p>L'invention concerne un procédé d'amplification des acides nucléiques dans lequel la détection des espèces amplifiées est augmentée par l'utilisation d'une amplification asymétrique. Une telle amplification est rendue asymétrique en utilisant des rapports divergents d'amorceurs d'amplification, ou en utilisant des amorceurs d'amplification non extensibles et/ou non clivables. La détection d'amplicons est améliorée du fait que le maintien des espèces à brins uniques d'amplicons facilite, durant l'amplification, leur capture directe par des sondes immobilisées sans avoir à recourir à des étapes de dénaturation.</p>		

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : C12Q 1/68	A1	(11) International Publication Number: <b>WO 00/61817</b> (43) International Publication Date: 19 October 2000 (19.10.00)
(21) International Application Number: PCT/US00/09742 (22) International Filing Date: 12 April 2000 (12.04.00)  (30) Priority Data: 09/290,452 12 April 1999 (12.04.99) US  (71) Applicant (for all designated States except US): NANOGEN/BECTON DICKINSON PARTNERSHIP [US/US]: 10398 Pacific Center Court, San Diego, CA 92121 (US).  (72) Inventors; and (75) Inventors/Applicants (for US only): EDMAN, Carl, F. [US/US]: 12545-C El Camino Real, San Diego, CA 92130 (US). NERENBURG, Michael, I. [US/US]: 8178 Gilman Court, La Jolla, CA 92037 (US). WESTIN, Lorelei, P. [US/US]: 8219 Finely Avenue, La Mesa, CA 91941 (US). CARRINO, John, J. [US/US]: 13129 Sea Knoll Court, San Diego, CA 92130 (US).  (74) Agents: MURDOCK, Douglas, C. et al.; Brobeck, Phleger & Harrison LLP, 12390 El Camino Real, San Diego, CA 92130 (US).		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: MULTIPLEX AMPLIFICATION AND SEPARATION OF NUCLEIC ACID SEQUENCES ON A BIOELECTRONIC MICROCHIP USING ASYMMETRIC STRUCTURES		
(57) Abstract  A method for amplifying nucleic acids is provided wherein detection of amplified species is enhanced by the use of asymmetric amplifications. Such amplification is made asymmetric by using divergent ratios of amplification primers or by using non-extending and/or non-cleavable amplification primers. Detection of the amplicons is improved because maintenance of single stranded species of amplicons during amplification facilitates their direct capture by immobilized probes without having to include denaturing steps.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**Description**

5

10

15

20

25

30

35

40

45

50

55



MULTIPLEX AMPLIFICATION AND SEPARATION OF NUCLEIC ACID  
SEQUENCES ON A BIOELECTRONIC MICROCHIP USING ASYMMETRIC  
STRUCTURES

FIELD OF THE INVENTION

This invention relates to devices, methods, and compositions of matter for performing active, multi-step, and multiplex nucleic acid sequence separation, amplification and diagnostic analyses. Generally, it relates to devices, methods, and compositions of matter for amplification and analysis of nucleic acid sequences in a sample. More specifically, the invention relates to methods, devices, and compositions of matter for amplifying and analyzing nucleic acids using novel strand displacement amplification technologies in combination with bioelectronic microchip technology. The devices and methods of the invention are useful in a variety of applications, including, for example, disease diagnostics (infectious and otherwise), genetic analyses, agricultural and environmental applications, drug discovery, pharmacogenomics, and food and/or water monitoring and analysis.

BACKGROUND OF THE INVENTION

The following description provides a summary of information relevant to the present invention. It is not an admission that any of the information provided herein is prior art to the presently claimed invention, nor that any of the publications specifically or implicitly referenced are prior art to that invention.

Definitions

The following descriptions of the inventions contained herein use numerous technical terms specific to the field of the invention. Generally, the meaning of these terms are known to those having skill in the art and are further described as follows:

As used herein, "sample" refers to a substance that is being assayed for the presence of one or more nucleic acids of interest. The nucleic acid or nucleic acids of interest may be present in a mixture of other nucleic acids. A sample, containing the nucleic acids of interest, may be obtained in numerous ways. It is envisioned that the

5 following could represent samples: cell lysates, purified genomic DNA, body fluids such as from a human or animal, clinical samples, food samples, *etc.*

10 As used herein, the phrases "target nucleic acid" and "target sequence" are used interchangeably. Both phrases refer to a nucleic acid sequence, the presence or absence of  
5 which is desired to be detected. Target nucleic acid can be single-stranded or double-stranded and, if it is double-stranded, it may be denatured to single-stranded form prior to  
15 its detection using methods, as described herein, or other well known methods. Additionally, the target nucleic acid may be nucleic acid in any form most notably DNA or RNA.

20 10 As used herein, "amplification" refers to the increase in the number of copies of a particular nucleic acid target of interest wherein said copies are also called "amplicons" or "amplification products".

25 As used herein, "amplification components" refers to the reaction materials such as enzymes, buffers, and nucleic acids necessary to perform an amplification reaction to form  
15 amplicons or amplification products of a target nucleic acid of interest.

30 As used herein, the phrase "multiplex amplification" refers to the amplification of more than one nucleic acid of interest. For example, it can refer to the amplification of multiple sequences from the same sample or the amplification of one of several sequences in a sample, as described in U. S. Patent Nos. 5,422,252 and 5,470,723 which are  
20 incorporated herein by reference. The phrase also refers to the amplification of one or more sequences present in multiple samples either simultaneously or in step-wise fashion.

35 As used herein, "oligonucleotide" refers to a molecule comprising two or more deoxyribonucleotides or ribonucleotides, preferably more than three. The length of an oligonucleotide will depend on how it is to be used. The oligonucleotide may be derived  
40 25 synthetically or by cloning. Oligonucleotides may also comprise protein nucleic acids (PNAs).

45 As used herein, "probe" refers to a known sequence of a nucleic acid that is capable of selectively binding to a target nucleic acid. More specifically, "probe" refers to an oligonucleotide designed to be sufficiently complementary to a sequence of one strand  
30 of a nucleic acid that is to be probed such that the probe and nucleic acid strand will hybridize under selected stringency conditions. Specific types of oligonucleotide probes  
50

are used in various embodiments of the invention. For example, a "ligation probe" describes one type of probe designed to bind to both a target nucleic acid of interest and to an amplification probe. Additionally, a "ligated probe" or a "ligated probe template" refers to the end product of a ligation reaction between a pair of ligation probes.

As used herein, the terms "primer molecule" and "primer" are used interchangeably. A primer is a nucleic acid molecule with a 3' terminus that is either "blocked" and cannot be covalently linked to additional nucleic acids or that is not blocked and possesses a chemical group at the 3' terminus that will allow extension of the nucleic acid chain such as catalyzed by a DNA polymerase or reverse transcriptase.

As used herein, the phrase "amplification primer" refers to an oligonucleotide primer used for amplification of a target nucleic acid sequence.

The phrase "primer extension," as used herein refers to the DNA polymerase induced extension of a nucleic acid chain from a free three-prime (3') hydroxy group thereby creating a strand of nucleic acid complementary to an opposing strand.

As used herein, the term "amplicon" refers to the product of an amplification reaction. An amplicon may contain amplified nucleic acids if both primers utilized hybridize to a target sequence. An amplicon may not contain amplified nucleic acids if one of the primers used does not hybridize to a target sequence. Thus, this term is used generically herein and does not imply the presence of amplified nucleic acids.

As used herein, "electronically addressable" refers to a capacity of a microchip to direct materials such as nucleic acids and enzymes and other amplification components from one position to another on the microchip by electronic biasing of the capture sites of the chip. "Electronic biasing" is intended to mean that the electronic charge at a capture site or another position on the microchip may be manipulated between a net positive and a net minus charge so that charged molecules in solution and in contact with the microchip may be directed toward or away from one position on the microchip or from one position to another position.

As used herein, the phrase "capture site" refers to a specific position on an electronically addressable microchip wherein electronic biasing is initiated and where molecules such as nucleic acid probes and target molecules are attracted or addressed by such biasing.

5

As used herein, the term "anchored" refers to the immobilization by binding of a molecule to a specified location on a microchip, such as a primer nucleic acid used in an SDA reaction, or a nucleic acid probe used to capture a target nucleic acid.

10

As used herein, the term "branched primer pair" refers to a pair of oligonucleotides that may be used as primers in an amplification reaction and which are connected together through a chemical moiety such that the oligonucleotides are susceptible to hybridization and use as amplification primers.

15

20

10

As used herein, the term "primer capture probes" refers to oligonucleotides that are used to hybridize to selected target nucleic acids and provide anchoring support for such nucleic acids to a capture site. Moreover, such oligonucleotides may function as amplification primers for amplifying said target nucleic acids.

25

15

As used herein, "hybridization" and "binding" are used interchangeably and refer to the non-covalent binding or "base pairing" of complementary nucleic acid sequences to one another. Whether or not a particular probe remains base paired with a polynucleotide sequence depends on the degree of complementarity, the length of the probe, and the stringency of the binding conditions. The higher the stringency, the higher must be the degree of complementarity, and/or the longer the probe for binding or base pairing to remain stable.

30

35

20

As used herein, "stringency" refers to the combination of conditions to which nucleic acids are subjected that cause double stranded nucleic acid to dissociate into component single strands such as pH extremes, high temperature, and salt concentration. The phrase "high stringency" refers to hybridization conditions that are sufficiently stringent or restrictive such that only specific base pairing will occur. The specificity should be sufficient to allow for the detection of unique sequences using an oligonucleotide probe or closely related sequence under standard Southern hybridization protocols (as described in *J. Mol. Biol.* 98:503 (1975)).

40

25

45

As used herein, "endonuclease" refers to enzymes (e.g., restriction endonucleases, etc.) that cut DNA at sites within the DNA molecule.

50

30

As used herein, a "restriction endonuclease recognition site" refers to a specific sequence of nucleotides in a double stranded DNA that is recognized and acted upon enzymatically by a DNA restriction endonuclease.

55

5  
10  
15  
As used herein, the term "nicking" refers to the cutting of a single strand of a double stranded nucleic acid by breaking the bond between two nucleotides such that the 5' nucleotide has a free 3' hydroxyl group and the 3' nucleotide has a 5' phosphate group. It is preferred that the nicking be accomplished with a restriction endonuclease and that this restriction endonuclease catalyze the nicking of double stranded nucleic acid at the proper location within the restriction endonuclease recognition site.

20  
25  
30  
As used herein, the phrase "modified nucleotide" refers to nucleotides or nucleotide triphosphates that differ in composition and/or structure from natural nucleotide and nucleotide triphosphates. It is preferred that the modified nucleotide or nucleotide triphosphates used herein are modified in such a way that, when the modifications are present on one strand of a double stranded nucleic acid where there is a restriction endonuclease recognition site, the modified nucleotide or nucleotide triphosphates protect the modified strand against cleavage by restriction enzymes. Thus, the presence of the modified nucleotides or nucleotide triphosphates encourages the nicking rather than the cleavage of the double stranded nucleic acid.

35  
40  
45  
As used herein, the phrase "DNA polymerase" refers to enzymes that are capable of incorporating nucleotides onto the 3' hydroxyl terminus of a nucleic acid in a 5' to 3' direction thereby synthesizing a nucleic acid sequence. Examples of DNA polymerases that can be used in accordance with the methods described herein include, *E. coli* DNA polymerase I, the large proteolytic fragment of *E. coli* DNA polymerase I, commonly known as "Klenow" polymerase, "Taq" polymerase, T7 polymerase, Bst DNA polymerase, T4 polymerase, T5 polymerase, reverse transcriptase, exo-BCA polymerase, etc.

50  
55  
As used herein, the term "displaced," refers to the removing of one molecule from close proximity with another molecule. In connection with double stranded oligonucleotides and/or nucleic acids, the term refers to the rendering of the double stranded nucleic acid molecule single stranded, i.e., one strand is displaced from the other strand. Displacement of one strand of a double-stranded nucleic acid can occur when a restriction endonuclease nicks the double stranded nucleic acid creating a free 3' hydroxy which is used by DNA polymerase to catalyze the synthesis of a new strand of nucleic acid. Alternatively, one nucleic acid may be displaced from another nucleic acid by the action of electronic biasing of an electrically addressable microchip.

As used herein, "ligase" refers to an enzyme that is capable of covalently linking the 3' hydroxyl group of a nucleotide to the 5' phosphate group of a second nucleotide. Examples of ligases include *E. coli* DNA ligase, T4 DNA ligase, *etc.*

As used herein, "ligating" refers to covalently attaching two nucleic acid molecules to form a single nucleic acid molecule. This is typically performed by treatment with a ligase, which catalyzes the formation of a phosphodiester bond between the 5' end of one sequence and the 3' end of the other. However, in the context of the invention, the term "ligating" is also intended to encompass other methods of connecting such sequences, *e.g.*, by chemical means.

The term "attaching" as used herein generally refers to connecting one entity to another. For example, oligomers and primers may be attached to the surface of a capture site. With respect to attaching mechanisms, methods contemplated include such attachment means as ligating, non-covalent bonding, binding of biotin moieties such as biotinylated primers, amplicons, and probes to streptavidin, *etc.*

As used herein, the term "adjacent" is used in reference to nucleic acid molecules that are in close proximity to one another. The term also refers to a sufficient proximity between two nucleic acid molecules to allow the 5' end of one nucleic acid that is brought into juxtaposition with the 3' end of a second nucleic acid so that they may be ligated by a ligase enzyme.

The term "allele specific" as used herein refers to detection, amplification or oligonucleotide hybridization of one allele of a gene without substantial detection, amplification or oligonucleotide hybridization of other alleles of the same gene.

As used herein, the term "three-prime" or "3'" refers to a specific orientation as related to a nucleic acid. Nucleic acids have a distinct chemical orientation such that their two ends are distinguished as either five-prime (5') or three-prime (3'). The 3' end of a nucleic acid contains a free hydroxyl group attached to the 3' carbon of the terminal pentose sugar. The 5' end of a nucleic acid contains a free hydroxyl or phosphate group attached to the 5' carbon of the terminal pentose sugar.

As used herein, the phrase "free three-prime (3') hydroxyl group," refers to the presence of a hydroxyl group located at the 3' terminus of a strand of nucleic acid. The phrase also refers to the fact that the free hydroxyl is functional such that it is able to

support new nucleic acid synthesis.

As used herein, the phrase "five-prime overhang" refers to a double-stranded nucleic acid molecule, which does not have blunt ends, such that the ends of the two strands are not coextensive, and such that the 5' end of one strand extends beyond the 3' end of the opposing complementary strand. It is possible for a linear nucleic acid molecule to have zero, one, or two, 5' overhangs. The significance of a 5' overhang is that it provides a region where a 3' hydroxyl group is present on one strand and a sequence of single stranded nucleic acid is present on the opposite strand. A DNA polymerase can synthesize a nucleic acid strand complementary to the single stranded portion of the nucleic acid beginning from the free 3' hydroxyl of the recessed strand.

As used herein, the term "bumper primer" refers to a primer used to displace primer extension products in SDA reaction. The bumper primer anneals to a target sequence upstream of the amplification primer such that extension of the bumper primer displaces the downstream amplification primer and its extension product.

As used herein, the terms "detected" and "detection" are used interchangeably and refer to the discernment of the presence or absence of a target nucleic acid or amplified nucleic acid products thereof.

As used herein, "label" refers to a chemical moiety that provides the ability to detect an amplification product. For example, a label on a nucleic acid may comprise a radioactive isotope such as  $^{32}\text{P}$  or non-radioactive molecule such as covalently or noncovalently attached chromophores, fluorescent moieties, enzymes, antigens, groups with specific reactivity, chemiluminescent moieties, and electrochemically detectable moieties.

The above definitions should not be understood to limit the scope of the invention. Rather, they should be used to interpret the language of the description and, where appropriate, the language of the claims. These terms may also be understood more fully in the context of the description of the invention. If a term is included in the description or the claims that is not defined above, or that cannot be interpreted based on its context, then it should be construed to have the same meaning as it is understood by those of skill in the art.

### Background art

Determining the nucleic acid sequence of genes is important in many situations. For example, numerous diseases are caused by or associated with a mutation in a gene sequence relative to the normal gene. Such mutation may involve the substitution of only one base for another, called a "point mutation." In some instances, point mutations can cause severe clinical manifestations of disease by encoding a change in the amino acid sequence of the protein for which the gene codes. For example, sickle cell anemia results from such a point mutation.

Other diseases are associated with increases or decreases in copy numbers of genes. Thus, determining not only the presence or absence of changes in a sequence is important but also the *quantity* of such sequences in a sample can be used in the diagnosis of disease or the determination of the risk of developing disease. Moreover, variations in gene sequences of both prokaryotic and eukaryotic organisms has proven invaluable to identifying sources of genetic material (*e.g.*, identifying one human from another or the source of DNA by restriction fragment length polymorphism (RFLP)).

Certain infections caused by microorganisms or viruses may also be diagnosed by the detection of nucleic acid sequences peculiar to the infectious organism. Detection of nucleic acid sequences derived from viruses, parasites, and other microorganisms is also important where the safety of various products is of concern, *e.g.*, in the medical field where donated blood, blood products, and organs, as well as the safety of food and water supplies are important to public health.

Thus, identification of specific nucleic acid sequences by the isolation of nucleic acids from a sample and detection of the sought for sequences, provides a mechanism whereby one can determine the presence of a disease, organism or individual. Generally, such detection is accomplished by using a synthesized nucleic acid "probe" sequence that is complementary in part to the target nucleic acid sequence of interest.

Although it is desirable to detect the presence of nucleic acids as described above, it is often the case that the sought for nucleic acid sequences are present in sample sources in extremely small numbers (*e.g.*, less than  $10^7$ ). The condition of small target molecule numbers causes a requirement that laboratory techniques be performed in order to amplify the numbers of the target sequences in order that they may be detected. There are many



5 well known methods of amplifying targeted sequences, such as the polymerase chain  
reaction (PCR), the ligase chain reaction (LCR), the strand displacement amplification  
10 (SDA), and the nucleic acid sequence-based amplification (NASBA), to name a few.  
These methods are described generally in the following references: (PCR) U.S. Patent Nos.  
5 4,683,195, 4,683,202, and 4,800,159; (LCR) EP Application No., 320,308 published June  
14, 1989; (SDA) U.S. Patent Nos. 5,270,184, and 5,455,166 and "Empirical Aspects of  
15 Strand Displacement Amplification" by G.T. Walker in *PCR Methods and Applications*,  
3(1):1-6 (1993), Cold Spring Harbor Laboratory Press; and (NASBA) "Nucleic Acid  
Sequence-Based Amplification (NASBA™)" by L. Malek *et al.*, Ch. 36 in *Methods in*  
20 10 *Molecular Biology*, Vol. 28: Protocols for Nucleic Acid Analysis by Nonradioactive  
Probes, 1994 Ed. P.G. Isaac, Humana Press, Inc., Totowa, NJ. (Each of the above  
references are hereby incorporated by reference.)

25 With respect to analyzing and/or identifying target nucleic acid amplified products,  
*i.e.*, "amplicons", other well known techniques have been typically used including  
15 comparative size, relative migration analyses (*e.g.*, Southern blot analysis) and  
hybridization analysis. However, comparative size or relative migration analyses are not  
30 optimal because they are undesirably slow and inaccurate. Additionally, while  
hybridization analysis offers many advantages over these methods, hybridization analysis  
is neither rapid nor sensitive as compared to the teachings of the present invention.

20 With respect to PCR technology, since thermal cycling is required, PCR is not  
35 optimal for use in a microelectronic environment because the heat fluctuations caused by  
the thermal cycling are detrimental to the capture sites located on the surface of a  
microchip. Thermal cycling gives rise to other problems as well including the requirement  
40 for complex instrumentation (*e.g.*, to ensure uniform heating, etc.), and, unacceptable time  
25 spans for completion of analysis (since each step must occur sequentially).

In contrast to PCR, the SDA technique is useful with microelectronic environments  
45 because it overcomes some of the above-described undesirable characteristics of PCR,  
*e.g.*, it is an isothermal method and the amplification process is asynchronous, and,  
therefore, more rapid. Although the use of SDA has advantages over PCR, SDA schemes  
30 as currently practiced typically include the use of solution-based amplification protocols  
50 (*e.g.*, disclosed in the above mentioned US Patent No. 5,455,166). Recent modifications

of the SDA technique have advanced the technique to minimizing the number of individually designed primers for amplification as described in US Patent No. 5,624,825. However, such advances do not benefit from enhancements realized in the current invention of electronically controlled hybridization.

Other amplification procedures include the use of probes that are bound to a solid support. However, such procedures have not provided a discernable advance in the art compared to the "anchored" SDA presented herein and performed in conjunction with an electronically addressable microchip. For example, US Patent No. 5,380,489 discloses a method for nucleic acid amplification and detection of target nucleic acids wherein an adhesive element is used to affix capture probes so that target molecules may be more easily captured and detected. This method does not address the issue of simultaneous amplification, capture, and detection as does the current invention. In another example, US Patent No. 5,474,895 discloses detection of nucleic acids using a polystyrene support-based sandwich assay. Again, such a method merely involves *passive* hybridization followed by subsequent detection following secondary *passive* hybridization of a probe.

Microchip arrays have also been used in association with nucleic acid amplification and detection. For example, miniaturized devices have been successfully developed for expression monitoring. See, e.g., M. Schena, *et al.*, 270 *Science* 467-470 (1995), M. Schena, *et al.*, 93 *Proc. Natl Acad. Sci. USA* 10614-619 (1996), J. DeRisi, *et al.*, 14 *Nat. Genet.* 457-60 (1996), R. A. Heller, *et al.*, 94 *Proc. Natl. Acad. Sci. USA* 2150-55 (1997), and J. DeRisi, *et al.*, 278 *Science* 680-86 (1997). Miniaturized devices have also been successfully developed for analysis of single nucleotide polymorphisms (SNPs) within an amplicon. See, e.g., Z. Guo, *et al.*, 15 *Nat. Biotechnol.* 331-35 (1997), and E. Southern, 12 *Trends Genet.* 110-15 (1996). (Each of the above publications are hereby incorporated by reference). These devices offer the potential for combining the specificity of hybridization with the speed and sensitivity of microchip technology. However, none have successfully provided a suitable miniaturized device for the present purposes.

For example, although micro-devices have been used to analyze multiple amplicons simultaneously (*i.e.*, multiplex analysis), such multiplex analysis has been possible only if hybridization conditions are compatible for each amplicon being tested. This detriment may be partially compensated for by careful capture probe design, by the

5 use of very long captures (e.g. cDNA for expression monitoring) (see, e.g., R. A. Heller, *et al.*, (1997) *supra*, and M. Schena, *et al.*, (1995) *supra*), or by extensive redundancy and  
10 overlap of shorter capture oligonucleotide sequences. However, taken together, these considerations have imposed limitations on the use of most microchip devices. Moreover,  
5 high levels of redundancy such as that used with short oligonucleotide capture sequences results in the requirement for large arrays and complex informatics programs to interpret  
15 data obtained, and still certain sequence-specific regions may remain difficult to analyze. Alternatively, the use of long capture oligonucleotides permits use of uniformly elevated hybridization temperatures. However, the use of long capture probes and elevated  
20 10 hybridization temperatures (e.g., in the range of 45 to 75°C) largely precludes single base pair mismatch analysis of highly related sequences.

Yet another disadvantage has become apparent with conventional microchips (e.g., those disclosed in US Patent Nos. 5,202,231 and 5,545,531, as well as in E. Southern *et al.*, *Genomics* 13, 1008-1017 (1992); M. Schena *et al.*, *Science* 279, 467-470 (1995); M.  
25 15 Chee *et al.*, *Science* 274, 610-614 (1996); and D. J. Lockhart *et al.*, *Nature Biotechnology* 14, 1675-1680 (1996) (all of which are herein incorporated by reference)), in that they depend upon passive hybridization and solution based amplification prior to the capture of amplified products on the microchips.

Further, many of these devices are unable to analyze and/or detect the amplification  
20 of target molecules from multiple samples simultaneously. In macroscopic devices, this latter problem is conventionally handled by "dot blot" formats in which individual samples occupy unique geometric positions with minimal contamination between samples. In  
35 contrast, for most microchips, the problem of detection and analysis usually requires expensive and complex nucleic acid deposition technology similar to dot blot macroscopic  
40 25 deposition but on a microscopic scale.

In another recent disclosure, (PCT WO96/01836), electronic microchips have been used in connection with PCR type amplification of nucleic acids. However, an  
45 amplification system requiring the simultaneous use of amplification enzymes and restriction enzymes for *increasing* the quantity of target amplicons at a specific capture site was not contemplated nor possible in that system. Rather, restriction digestion of captured  
30 50 nucleic acid species was considered in connection with the removal of double stranded

nucleic acid species from capture sites following PCR type amplification procedures with detection of target species occurring subsequent to enzymatic cleavage. Moreover, that system provided anchored amplification primers complementary to only one strand of a target nucleic acid that were functional in a PCR reaction.

Like other microchip based amplification and detection platforms, the invention conceptualized in the PCT WO 96/01836 publication is substantially limited as compared to the SDA on electronically addressable microchips disclosed herein because the PCR type amplification of target species as taught in that publication required repeated disruption of double stranded species as well as functionality of solution based reverse primers. Such a situation results in the reduction of efficient amplification due to primer-primer interactions while use of restriction enzymes is inhibited due to fluctuations in reaction buffer conditions.

Finally, other aspects of amplification and detection of nucleic acids have been problematic and /or not optimal. One such problem has been the loss of specificity in the restriction endonuclease cleavage of nucleic acids by restriction enzymes. For example, it is known that some restriction endonucleases lose specificity for their DNA recognition sequence with increased osmotic pressure or reduced water activity. C. R. Robinson *et al.* *J. Mol. Biol.* 234: 302-306 (1993). With reduced water activity, the restriction endonucleases will cleave DNA at recognition sites that differ by one base pair from the normal recognition site. The restriction sites that are off by one base pair are called "star" sites and the endonucleases recognition and cleavage of these star sites is called "star activity."

Robinson *et al.* found that bound water participates in sequence specificity of *EcoRI* DNA cleavage (*Biochemistry* 33(13):3787-3793(1994)), and further found that increasing hydrostatic pressure by conducting the reactions at elevated pressure from 0 to 100 atm. inhibited and ultimately eliminated star activity induced by osmotic pressure for *EcoRI*, *PvuII*, and *BamHI*, but not for *EcoRV*. (*Proc. Natl. Acad. Sci. USA* 92:3444-3448 (1995)). One recurrent problem with SDA that relies on restriction endonucleases is the frequency with which non-target sequences are amplified in a primer-independent manner due to star activity. Thus, there is a need to reduce or eliminate star activity in SDA reactions. In one embodiment of the current invention, we provide for the elimination of

5 such star activity in SDA reactions by application of a high pressure SDA method.

10 In addition to advancing SDA technology by eliminating star activity, we also  
provide for various other advancements in the detection of nucleic acids using SDA in  
combination with a bioelectronic microchip. For example, amplification and separation of  
5 nucleic acid sequences may be carried out using ligation-dependent SDA. In contrast to  
ligation-dependent amplification procedures known in the art that require the amplified  
15 products to be separated from the starting material by a capture step, or that require that  
free ligation probe be separated from bound probe prior to ligation, the current invention  
eliminates the need to make separate isolation steps. Further, the current invention  
20 improves upon the SDA amplification process by eliminating the need for bumper primers  
as they have been used in the art. For example, typical ligation-dependent amplification  
procedures include capture steps by labeling one of the primers used during amplification.  
25 Separation may occur prior to ligation to prevent template independent ligation of the  
primers or separation may occur following ligation to isolate target DNA amplicons from  
15 the non-labeled/amplified DNA. Target DNA amplicons containing this label are separated  
from the non-labelled/amplified DNA. This separation requires an extra step following  
amplification. This extra manipulation of the sample increases the complexity of the  
30 procedure and thereby renders it less useful as a simple alternative to other current DNA  
amplification methods such as PCR. This extra manipulation of sample also hinders  
20 automation of the amplification procedure. In one embodiment of the current invention a  
ligation-dependent SDA method is provided that eliminates such extra steps facilitating  
35 automation of amplification and detection of target nucleic acids.

40 In other embodiments, we have provided additional advancements in nucleic acid  
amplification and detection technology using SDA and electronically addressable  
25 microchips which advancements collectively show that a need remains for devices,  
methods, and compositions of matter for efficiently and optimally amplifying, detecting  
and analyzing target nucleic acid sequences of interest.

5

10

15

20

25

30

35

40

45

50

55

## SUMMARY OF THE INVENTION

This invention relates broadly to devices, methods, and compositions of matter for the multiplex amplification, detection and analysis of nucleic acid sequences wherein the amplification, detection and analysis is optimally accomplished using SDA in combination with bioelectronic microchip technology. The invention provides various efficient and optimal methods of amplifying target nucleic acids of interest as well as methods for analyzing resultant amplicons. In addition, the invention enables the amplification and analysis (either sequentially or simultaneously) of multiple samples containing target nucleic acids on a single open bioelectronic microchip.

In one aspect of this invention, the microchip device is an electronically controlled microelectrode array. *See*, PCT application WO96/01836, the disclosure of which is hereby incorporated by reference. In contrast to the passive hybridization environment of most other microchip devices, the electronic microchip devices (or active microarray devices) of the present invention offer the ability to actively transport or electronically address nucleic acids to discrete locations on the surface of the microelectrode array, and to bind the addressed nucleic acid at those locations to either the surface of the microchip at specified locations designated "capture sites" or to nucleic acids previously bound at those sites. *See*, R. Sosnowski, *et al.*, 94 *Proc. Natl. Acad. Sci. USA* 119-123 (1997), and C. Edman, *et al.*, 25 *Nucleic Acids Res.* 4907-14 (1997). The use of these active microarrays circumvent many of the limitations encountered by passive microdevices.

The active microchip arrays of the present invention overcome the size dependency of capture oligonucleotides and the complexity requirements of passive microdevices. Also, the microchip arrays of the present invention allow multiple independent sample analyses upon the same open microarray surface by selectively and independently targeting different samples containing nucleic acids of interest to various microelectrode locations. In other words, they allow parallel multiple sample processing on an open array. As mentioned above, traditional nucleic acid detection methodologies are restricted by the frequently long amplification and hybridization times required to achieve resolvable signals. An additional limitation to such methodologies is the inability to carry out multiplex hybridization events upon their analytical surfaces, thereby restricting information obtainable in any one assay. Both of these limitations are overcome in the

5 present invention by use of active microelectronic arrays capable of selectively targeting  
and concentrating DNA to specific sites on the array. A further strength of these devices is  
10 the power to perform electronic hybridization and denaturation to discriminate single base  
polymorphisms. Thus, these active microelectrode arrays demonstrate the flexibility to  
5 handle a wide variety of tasks upon a common platform, beyond those seen with other  
microdevices.

15 The present invention preferably uses an amplification method different from  
traditional PCR. Specifically, the present invention uses strand displacement amplification  
(SDA). SDA is an amplification methodology that has sufficient sensitivity and  
20 10 robustness to rapidly (*e.g.*, in about 15-45 minutes) and exponentially amplify a small  
number of target molecules over a complex background. *See, e.g.*, C. Spargo, *et al.*, 10  
*Molecular and Cellular Probes* 247-56 (1996). In contrast to PCR, SDA is an isothermal  
25 technique that requires simpler thermal control and associated instrumentation. SDA is  
more compatible with a unified amplification-hybridization-detection system (*i.e.* a system  
15 wherein all steps are unified in one place, *e.g.*, on a microarray chip) for rapid analyses of  
nucleic acids. This is primarily due to the fact that SDA does not require conditions (*e.g.*  
30 thermal cycling) which could be detrimental to the microarray of an electronically  
addressable microchip.

35 The efficiency of amplification reactions in *passive* hybridization wherein probes  
20 designed to capture target and amplicon nucleic acid molecules are anchored to the surface  
of the microarray is limited during the initial phases of amplification due to the low  
frequency of hybridization of target nucleic acid species to the appropriate primers located  
40 on the tethering surface. Typically, this process requires hours, even in reduced volumes  
of solution. However, the efficiency of this process is dramatically increased by  
25 electronically concentrating, (*i.e.* addressing), the nucleic acid to the vicinity of "anchored"  
primers, thereby increasing the frequency of encounter between the solution phase target  
nucleic acid and the anchored primers. Whereas prior concepts used PCR in connection  
45 with only one of the two amplification primers necessary for PCR amplification anchored  
to a specific site on the microarray, the current invention contemplates that both  
30 amplification primers necessary for SDA are anchored to a specific capture site on the  
microarray. Thus, in one embodiment of the invention, electronically concentrating and  
50

5 hybridizing the target nucleic acid to the surface of a microchip (*i.e.*, capture sites) prior to  
the introduction of amplification reaction buffers, enzymes, nucleotides, *etc.*, benefits  
10 greatly "anchored" amplification reactions, such as "anchored SDA", as described below.  
The rapid concentration and subsequent specific hybridization of template nucleic acid  
5 molecules to their complementary anchored amplification primers permits the surface of  
the array to be washed, removing unwanted and possibly interfering non-target nucleic  
15 acid from the reaction environment.

Employing electronic addressing of target nucleic acids to specific locations on the  
microarray has at least three other advantages over prior passive hybridization  
20 technologies. First, the overall time and efficiency of the amplification process is  
dramatically improved since a major rate-limiting step (that of the time required for the  
template to find the anchored primers) is removed from the overall reaction rate. Also, the  
25 use of electronic addressing acts to electronically concentrate target nucleic acids such that  
hybridization of the target species to the anchored amplification probes increases the  
15 number of target molecules at the selected site as compared to the number of target  
molecules that would be found at any particular site on a non-electronic, passive  
hybridization microarray for an equivalent time period. The result is that the absolute  
30 numbers of starting molecules for the amplification process is dramatically increased  
resulting in improvement in both the overall yield of amplification products and the  
20 sensitivity to lower starting template numbers.

The second advantage is that discrete target nucleic acids can be applied to specific  
locations upon the array surface thereby allowing multiple, different nucleic acid samples  
to be simultaneously amplified on one array. Alternatively, a nucleic acid sample can be  
40 targeted to several different locations, each containing specific sets of amplification  
25 primers so that multiple different amplification reactions can be simultaneously carried out  
from a single sample. As noted above, the ability to remove unnecessary and unhybridized  
DNA from the reaction mixture significantly aids this process.

45 A third advantage to this approach is that following an amplification reaction, the  
amplicons which have been addressed and bound to a specific site on the array are then  
30 available in a site-specific fashion for subsequent analyses, such as by (1) the introduction  
of fluorescently labeled nucleotides or (2) the hybridization of oligonucleotides at the end  
50



5 of the reaction by denaturation of the amplified material followed by hybridization with an appropriate reporter oligonucleotide having specificity for the tethered amplicon.

10 As is described herein, the ability of electronic targeting used in connection with the combination of an electronically addressable microchip and SDA to overcome the  
5 above-described limitations of prior methods is demonstrated in two examples of amplicon analysis. First, as described in more detail below, use of a common highly conserved  
15 locus (e.g., 16S rRNA) which is shared by numerous species of bacteria may be applied to multiple comparative analyses of individual samples to identify different bacteria types. Second, also described in more detail below, the electronic microarray of the present  
20 10 invention is used to simultaneously analyze multiple individual patient samples for the presence of the human Factor V Leiden (R506Q) gene mutation. The human Factor V Leiden (R506Q) gene indicates a predisposition to activated protein C resistance and venous thrombosis. This example shows successful parallel sample analyses from  
25 multiple patients. The test material used in this multiple patient sample analysis provides  
15 another aspect of the present invention, namely, an allele-specific amplification method using SDA, also described in more detail below.

30 Other aspects of the present invention are directed to various new amplification methods. Such novel SDA methods of the present invention are useful for providing amplicons for various analyses. For example, some of the SDA methods described herein  
20 are useful to optimize amplification conditions for conducting amplification on an electronically addressable microchip array. Other SDA methods are useful to provide  
35 amplicons particularly suited for use on an electronically addressable microchip array. Still other SDA methods are useful to optimize analysis conditions for an analysis  
40 conducted on an electronically addressable microchip array.

25 One embodiment of a SDA method of the present invention, more specifically, comprises an allele-specific SDA method. The method preferably selectively amplifies  
45 only those strands that include a specific allele. The method preferably uses amplifying primers designed so their 3' terminus complements the nucleotide sequence of the desired allele. The primer may also preferably include a biotin moiety on its 5' end to provide a  
30 facile mechanism for capturing the amplicon and/or target nucleic acid onto a capture site  
50 either prior to amplification or after amplification following electronic targeting.

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

Additionally, in another allele-specific embodiment, a method is provided for analyzing multiple samples containing nucleic acids for the presence of alleles of a given gene, which comprises amplifying the nucleic acids in each sample by "two-strand" SDA to produce amplicons, wherein the first amplification uses primers specific for a first allele and the second amplification uses primers specific for a second allele, electronically addressing the amplicons on a microarray, hybridizing one or more reporter probes to the bound amplicons, and detecting the presence and location of the reporter probes on the microarray.

10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In another embodiment of the current invention, a unique combination of SDA and simultaneous detection of amplification products on an electronically addressable microchip is provided. In a preferred embodiment, SDA is carried out at the surface of a designated position on an electronic microchip wherein both upstream and downstream primers necessary for amplification are anchored to the same discrete capture site on a microarray. In one such embodiment, the primers are paired using a unique branched moiety that is "anchored" to the surface of the microchip. This branched primer pair design provides closely spaced primers having a defined distance and location from one another. This arrangement further provides a means by which the rate of SDA can be controlled. Moreover, combined with other elements of the invention, single stranded amplification products being created at the location of the primer pair may be easily and quickly addressed and captured by unused branched primer pairs onto the same or adjacent designated capture sites on the electronic microchip for further SDA.

10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In a preferred embodiment, each primer of the above mentioned primer pair further includes nucleic acid sequence encoding one strand of an endonuclease restriction site positioned 5' to a nucleic acid sequence having nucleic acid sequence complementary with a target molecule. In a further preferred embodiment, the sequence of the restriction sites in the primers are *unmodified* in that the nucleic acid backbone comprises a natural phosphate backbone that is cleavable by action of the restriction enzyme. Additionally, the restriction sites useful in SDA may be any restriction site typically used in SDA procedures as disclosed in the references incorporated herein such as HincII, HindII, BsoBI, Aval, Fnu4HI, Tth111I, and NciI. Other endonucleases can also be used in this method including BstXI, BsmI, BsrI, BsaI, NlaV, NspI, PfiMI, HphI, AlwI, FokI, AccI, TthIII,

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

Mra I, Mwo I, Bsr BI, Bst NI, Bst OI, and Bsm AI. Additionally, the enzyme need not be thermophilic. Moreover, it is a further preferred embodiment that the double stranded SDA amplification product produced during primer extension become hemimethylated or hemiphosphorothiolated (or other hemimodified form known to those skilled in the art) so that the double stranded SDA amplification product can be properly "nicked" at the primer restriction site for normal SDA amplification. For example, the substituted deoxynucleosidetriphosphate should be modified such that it will inhibit cleavage in the strand containing the substituted deoxynucleotides but will not inhibit cleavage on the other strand. Examples of such substituted deoxynucleosidetriphosphates include 2'deoxyadenosine 5'-O-(1-thiotriphosphate), 5-methyldeoxycytidine 5'-triphosphate, 2'-deoxyuridine 5'-triphosphate, and 7-deaza-2'-deoxyguanosine 5'-triphosphate.

In an alternative preferred embodiment, a restriction site may be used in the SDA procedure that does not require the nucleic acid backbone of the restriction site to be modified as described above. For example, BstNBI may be used in connection with its restriction site to nick the nucleic acid as it does not require modification to achieve single stranded nicks. Instead, BstNBI performs single stranded nicks as a natural activity.

The nucleic acid segments of the primer pair complementary to target sequence may be any useful length that will allow hybridization under temperature and buffer conditions appropriate for proper function of SDA on the microchip. Typically, the target sequences of the primer pair have sequence that is complementary with portions of target nucleic acids that are spaced anywhere from 60 to 120 bases upstream or downstream, as the case may be, from one another. In all cases each primer of the primer pair is complementary to different strands (*i.e.*, the plus strand or the minus strand) of the target sequence. Additionally, where the primer pair is on a branched moiety the spacing between the primers on the branched connecting moiety may be adjusted by molecular spacer elements to optimally enhance the efficiency of the SDA reaction. Such spacer elements may comprise polyethylene glycol polymers, polyamino acids, or nucleic acids.

In another preferred embodiment, the spaced primers may be attached to a branched molecular structure (*e.g.*, a 'Y' shaped structure) at their respective 5' termini. The branched structure is itself then anchored via a free branch of the Y to designated capture pad sites on the microchip. Attachment chemistry to the microchip surface may be

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

by streptavidin/biotin coupling well known in the art. Alternatively, attachment chemistry may include chemistry comparable to that disclosed in any of US Patent Nos. 5,668,258, 5,668,257, 5,677,431, 5,648,470, 5,623,055, 5,594,151, and 5,594,111, herein incorporated by reference. In one preferred embodiment, the branched molecules are formed by nucleic acids attached to an amino acid. In another alternate embodiment, the branched molecules are formed by adding different spacers, such as polyethylene glycol polymers, polyamino acids, or nucleic acids between the nucleic acid primers and a bifunctionally branched amino acid (e.g. lysine).

In yet another embodiment, the anchored SDA amplification primers need not be branched but instead merely anchored individually to the capture site in close proximity to each other. Attachment chemistry may be accomplished as described above.

In another preferred aspect of the invention, amplification of target nucleic acids is carried out exclusively at the site of an anchored primer pair thereby avoiding the uncertainties of amplification rate commonly associated with solution-based amplification. Particularly, as compared with solution-based amplification, the amplification of multiple targets or multiplex amplification is markedly improved. It is probable that such improvement is due to the avoidance of competition between primers and/or avoidance of primer-primer interactions that may inhibit binding to target sites. Amplification is kept at one location by the combined influence of electronic addressing of target molecules and SDA products to capture pad SDA sites and by the fact that the primers that allow amplification (*i.e.*, the branched or unbranched primer pairs) are anchored to a fixed location.

In another preferred aspect of the invention, the target nucleic acid is electronically addressed to the specific site on the microchip prior to amplification. This aspect is an advance over passive hybridization technology in several ways. First, since nucleic acids in a sample solution containing target nucleic acid species are electronically addressed to specific sites on the microchip, the target molecules have a preferred advantage of contacting the primer pair designed to capture the target molecule. Secondly, in the event single stranded nucleic acid target molecules must be generated, conditions in the sample solution that allow for formation of single stranded species must only be accomplished once rather than repeatedly as is normally the case with PCR and solution-based

5 amplification. Third, the electronic addressing and annealing of the target species to  
specific capture sites on the chip may be carried out in low salt conditions, a situation that  
10 is markedly in contrast to classical passive hybridization technology. Low salt conditions  
(and electronic addressing) enhance the hybridization of single stranded target species to  
5 capture primers because such conditions help reduce the reannealing of target nucleic acid  
strands to their respective complementary strands.

15 In another preferred embodiment, the anchored SDA methods of the current  
invention provide improved efficiency because only one target specific "bumper" primer is  
required for annealing to the target molecule at a position on the target 5' to the target  
20 annealing position of one or the other anchored primers. In another embodiment, two  
bumper primers may be included (as in traditional SDA) but inclusion of two primers is  
not necessary. Rather, the use of two bumper primers only facilitates initiation of priming  
25 from either direction on any one pair of primer capture probes depending upon which of  
the two strands of target nucleic acid are first captured by the branched primer pair.  
15 Inclusion of two bumper primers may further enhance the rate of amplicon formation.

30 In yet another aspect of this invention, a method of amplification of a target nucleic  
acid sequence (and its complementary strand) in a sample using SDA under elevated  
pressure is provided. By elevating the pressure, the efficiency of the amplification is  
enhanced because the specificity of the restriction endonuclease for its target sequence is  
20 increased. The method involves the steps of 1) isolating nucleic acids suspected of  
35 containing the target sequence from a sample, 2) generating single stranded fragments of  
target sequences, 3) adding a mixture comprising (a) a nucleic acid polymerase, (b)  
deoxynucleosidetriphosphates, a phosphorothioated dNTP, endonuclease, and (c) at least  
40 one primer which (i) is complementary to a region sometimes at or along a portion of the  
25 target near the 3' end of a target fragment, and (ii) further has a sequence at its 5' end  
which is a recognition sequence for a restriction endonuclease, and 4) allowing the mixture  
to react under elevated pressure for a time sufficient to generate amplification products.  
45 Where the target nucleic acid fragments comprise double stranded nucleic acids, the  
method further comprises denaturing the nucleic acid fragments to form single stranded  
30 target sequences. Where the nucleic acids comprise RNA, it is preferable to first use  
reverse transcriptase in order to convert RNA to DNA, however, RNA is specifically  
50

5 included in all embodiments of the invention.

10 In a further embodiment, a method of SDA in conjunction with an electronic microchip is provided wherein the SDA reaction is ligation-based. In this embodiment, two sets of primers are used wherein one primer set is designed so that the primers bind to  
5 one strand of a target sequence adjacent to one another while each of the primers of the second set are designed to bind to a portion of one of the primers of the first primer set  
15 while the other of the second primer set is complementary to a portion of the other of the first primer set (*i.e.*, same as the target strand sequence). When this embodiment is used, it will be apparent that SDA may be accomplished without the involvement of bumper primers. In a preferred embodiment, one of the two primer sets may be "anchored" as  
20 described herein.

25 In another embodiment, a method of ligation-based SDA is provided where the method is unassisted by an electronic microchip. In this embodiment it is not necessary to, *inter alia*, anchor any primers, which is a procedure that assists in separating primer sets  
15 during multiplex amplification, because the primers are universal -- there is no need to direct target sequences to the 'correct' primers.

30 In a particular embodiment of the ligation-based SDA method, the probe set designed to anneal to a target sequence must become ligated to form a "ligated probe template" which template is capable of supporting SDA. In a further preferred  
20 embodiment, the ligation-based reaction uses a single pair of amplification primers (*i.e.*,  
35 the second primer set mentioned above) which are universally applicable to amplification of all target molecules in a multiplex test providing in turn for decreased non-target amplification as well as decreased primer competition interactions due to the absence of  
40 bumper primers.

25 In a further preferred embodiment, the ligated probe template is modified so that it can not be extended from its 3' end during initial SDA reaction steps. Modifying the relevant ligation probe prevents the formation of a double stranded nucleic acid the 3' end  
45 of which may be cleaved by restriction endonuclease due to formation of what would be a cleavable restriction site, as explained in more detail below. This modification also  
30 prevents amplification of ligated probe template that may result from the target-sequence-independent ligation of the ligation probes.  
50  
55

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In another preferred embodiment of the ligation-based SDA method, the pair of probes used to target a nucleic acid of interest and create a ligation probe template are bifunctional in that each probe of the pair contains a target binding sequence and an "amplification primer" binding sequence (*i.e.*, the second primer set mentioned above). The sequences specific for target binding are chosen so that they are complementary to adjacent sequences of target DNA. The portions of the ligation probe template primers having nucleic acid sequence used in amplification are chosen so that a single set of amplification primers can be used for all target species of interest during SDA.

10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In a further embodiment, a first amplification primer binds to the ligated probe template at the 3' end of the ligated probe template such that there is created two 5' overhangs. *See* Figure 23(a). Double stranded nucleic acids with 5' overhangs are normally capable of supporting nucleic acid synthesis from the 3' end of the recessed strand by a DNA polymerase. As is well known in the art, DNA polymerase functions by extending the length of one strand of a nucleic acid by incorporating bases to the strand that are complementary to the opposing strand.

10  
15  
20  
25  
30  
35  
40  
45  
50  
55

However, in a further preferred embodiment, nucleic acid synthesis from the 3' terminus of the ligated probe template is prevented due to the 3' terminus having a modification to keep it from extending. Those in the art understand that this modification may take many forms including but not limited to: creating a 3' base mismatch between the ligated probe and the amplification primer; using a 3' terminal dideoxy nucleotide; or modifying the chemical moiety present at the 3' carbon of the pentose sugar of the nucleic acid backbone by, for example, replacing the free 3'hydroxyl group with a phosphate group, a biotin moiety, or by adding other blocking groups which are well known to those in the art. (*See* U.S. Patents 5,516,663 and 5,573,907 and 5,792,607, incorporated herein by reference, discussing various reagents that can be used to modify ends of the ligation probes to prevent target independent ligation). This modification prevents the formation of a double stranded nucleic acid which could be improperly "nicked" by endonuclease during the ligation-based amplification process. This modification also prevents amplification of ligated probe template that may result from the target sequence independent ligation of the ligation probes and prevents 3' extension when ligated probe is bound to primer. This modification also allows the ligation and amplification reactions to

5 proceed without an additional capture step.

10 In a further preferred embodiment, the ligation probes are designed to include sequences encoding endonuclease restriction sites, such that these sites are located near the 5' and 3' ends of the ligated probe template. Restriction endonuclease present in the  
5 reaction mixture may nick the double stranded nucleic acid so that SDA may proceed. Nicking of the DNA rather than cleavage occurs because the strand complementary to the  
15 5' end of the ligated probe is synthesized during SDA using nucleotides that include a modified nucleotide (for example dATP $\alpha$ S, or dCTP $\alpha$ S).

20 In a further embodiment, the amplicons arising from ligation-based SDA may be addressed to capture sites following their respective formation (whether their amplification is made to occur by SDA in solution or directly on the capture sites by primers that are addressed to the capture sites prior to amplification as described herein).

25 In yet another embodiment of the invention, several means by which the presence of target nucleic acids in a sample may be detected are available due to the combined application of the electronic addressable chip and anchored SDA. For example, in a  
15 preferred embodiment, amplicons that are addressed to capture sites may be discerned directly by fluorescence, *i.e.*, a fluorochrome may be included in the buffer so that detection is simultaneous with the production of amplicons. Examples of such fluorescing  
30 compounds include Bodipy-derivatives, Cy-derivatives, fluorescein-derivatives, and rhodamine-derivatives all of which are well known in the art. Alternatively, detection of  
20 nucleic acids at capture sites may be carried out directly using chemiluminescence or electrochemiluminescence. Chemiluminescence incorporates the use of an enzyme linked to a reporter oligonucleotide which, when activated with an appropriate substrate, emits a  
35 luminescent signal. Examples of such enzymes include horseradish peroxidase and alkaline phosphatase both of which are well known in the art. Electrochemiluminescence  
40 (ECL) is a highly sensitive process (200 fmol/L) with a dynamic range of over six orders of magnitude. In this system, reactive species are generated from stable precursors at the  
25 surface of an electrode. These precursors react with each other to form the excited state of the label attached to the DNA strand. The excited state decays to the ground state through  
30 a normal fluorescence mechanism, emitting a photon having a wavelength of 620 nm.



5

10

15

20

25

30

35

40

45

55

The amplification products generated using the primers disclosed herein may also be detected by a characteristic size, for example, on polyacrylamide or agarose gels stained with ethidium bromide. Alternatively, amplified target sequences may be detected by means of an assay probe, which is an oligonucleotide tagged with a detectable label. In one embodiment, at least one tagged assay probe may be used for detection of amplified target sequences by hybridization (a detector probe), by hybridization and extension as described by Walker, *et al.* (1992, *Nucl. Acids Res.* 20:1691-1696) (a detector primer) or by hybridization, extension and conversion to double stranded form as described in EP 0678582 (a signal primer). Preferably, the assay probe is selected to hybridize to a sequence in the target that is between the amplification primers, *i.e.*, it should be an internal assay probe. Alternatively, an amplification primer or the target binding sequence thereof may be used as the assay probe.

The detectable label of the assay probe is a moiety which can be detected either directly or indirectly as an indication of the presence of the target nucleic acid. For direct detection of the label, assay probes may be tagged with a radioisotope and detected by autoradiography or tagged with a fluorescent moiety and detected by fluorescence as is known in the art. Alternatively, the assay probes may be indirectly detected by tagging with a label that requires additional reagents to render it detectable. Indirectly detectable labels include, for example, chemiluminescent agents, enzymes which produce visible reaction products and ligands (*e.g.*, haptens, antibodies or antigens) which may be detected by binding to labeled specific binding partners (*e.g.*, antibodies or antigen/haptens). Ligands are also useful immobilizing the ligand-labeled oligonucleotide (the capture probe) on a solid phase to facilitate its detection. Particularly useful labels include biotin (detectable by binding to labeled avidin or streptavidin) and enzymes such as horseradish peroxidase or alkaline phosphatase (detectable by addition of enzyme substrates to produce colored reaction products). Methods for adding such labels to, or including such labels in, oligonucleotides are well known in the art and any of these methods are suitable for use in the present invention.

Examples of specific detection methods that may be employed include a chemiluminescent method in which amplified products are detected using a biotinylated capture probe and an enzyme-conjugated detector probe as described in U.S. Patent No.

5,470,723. After hybridization of these two assay probes to different sites in the assay region of the target sequence (between the binding sites of the two amplification primers), the complex is captured on a streptavidin-coated microtiter plate by means of the capture probe, and the chemiluminescent signal is developed and read in a luminometer. As another alternative for detection of amplification products, a signal primer as described in EP 0678582 may be included in the SDA reaction. In this embodiment, labeled secondary amplification products are generated during SDA in a target amplification-dependent manner and may be detected as an indication of target amplification by means of the associated label.

In another alternative detection method, a target specific primer, (*i.e.*, a target signal primer which is a primer that is not a bumper primer or an anchored primer), designed to anneal to the target sequence at a position other than at the anchored primer or bumper primer sites may be included in the amplification step procedure. This signal primer may be labeled with a signal molecule that may in turn be used to detect an extension product formed from extension of the signal primer during SDA. For example, such label may comprise biotin that may be captured to a microchip location containing streptavidin which capture may be detected by presence of a fluorochrome.

In still another aspect of the invention, use of a signal primer elongation product or amplicon provides for a means by which the molar ratio of one target amplicon strand over the other may be produced so that single stranded amplified species of the target sequence may be maintained for capture by capture probes located at specific sites on the microchip. In other words, the signal primer allows "asymmetric SDA". Moreover, the amplified signal primed amplicons may be electronically addressed to secondary capture sites which facilitates further reduction in background signal for enhanced detection.

For commercial convenience, amplification primers for specific detection and identification of nucleic acids may be packaged in the form of a *kit*. Typically, such a kit contains at least one pair of amplification primers. Reagents for performing a nucleic acid amplification reaction may also be included with the target-specific amplification primers, for example, buffers, additional primers, nucleotide triphosphates, enzymes, *etc.* The components of the kit are packaged together in a common container, optionally including instructions for performing a specific embodiment of the inventive methods. Other

optional components may also be included in the kit, e.g., an oligonucleotide tagged with a label suitable for use as an assay probe, and/or reagents or means for detecting the label.

#### BRIEF DESCRIPTION OF THE DRAWINGS

The file of this patent contains at least one drawing executed in color. Copies of this patent with color drawing(s) will be provided by the Patent and Trademark Office upon request and payment of the necessary fee.

Figure 1A shows a cross-sectional view of an embodiment of the bioelectronic chip of the present invention.

Figure 1B shows a perspective view of the bioelectronic chip from Fig. 1A.

Figure 2A shows a schematic representation of a bacterial 16S rRNA gene comprising a divergent region (having a different sequence per bacterial strain) flanked on both sides by conserved regions (having the same sequence in each bacterial strain). BBs and Bba represent bacterial sense and antisense bumper primers respectively. Bas and Baa represent bacterial sense and antisense amplification primers respectively. Identification of the bacterial strains tested are in the sequence listing.

Figure 2B shows the results of 16S rRNA encoding SDA amplification products resolved on a 1% agarose gel stained with ethidium bromide showing specific amplification of the divergent regions from each strain.

Figure 2C shows one aspect of a sandwich assay format used for nucleic acid hybridization on microarrays of the present invention wherein the assay format utilizes a universal capture probe and a sequence specific reporter.

Figure 2D shows a sandwich assay format used for nucleic acid hybridization on microarrays of the present invention wherein the assay format utilizes a sequence specific capture probe and a universal reporter.

Figure 3A shows *Salmonella*-specific BTR labeled reporter used for *passive* hybridization of SDA amplicons on a microarray wherein the capture sites of the microarray include as a control for non-specific binding of the reporter oligonucleotide to the capture probes or permeation layer itself a site containing capture probes but no target (+C/-T) and a site containing no capture probe or target (C/-T).

Figure 3B shows a comparison of the relative fluorescence observed for each

bacteria when SDA amplicons were generated and *electronically* addressed to individual sites on a microarray using *universal* capture probes, and sequence-specific btr-labeled reporter probes (designed in the divergent region of the 16S rRNA gene) were passively hybridized to discriminate various bacterial strains.

Figure 3C shows a comparison of the relative fluorescence observed for each bacteria when SDA amplicons were generated and *electronically* addressed to individual sites on a microarray using *sequence-specific* capture probes, and universal btr-labeled reporter probes (designed in the conserved region of the 16S rRNA gene) were passively hybridized to the captured material.

Figure 4A shows a polyacrylamide gel analysis of the allele-specific reactions from five patient samples analyzing for Factor V Leiden mutation in each wherein each genomic DNA sample was amplified twice with allele-specific SDA using either the normal genotype (Factor V R506), W, or the Leiden mutation (Factor V Q506), M.

Figure 4B shows a histogram comparing the fluorescence present at each addressed site on the array of the allele-specific reactions from three of the five patient samples of Figure 4A.

Figure 5A shows a diagram of a first scheme of incorporating a fluorescent species in an amplification reaction for detection purposes.

Figure 5B shows a diagram of a second scheme of incorporating a fluorescent species in an amplification reaction for detection purposes.

Figure 6A shows a fluoroscopic analysis of a microchip where the SDA template was absent as a control.

Figure 6B shows a fluoroscopic analysis of a microchip where BsoBI was not included in the reaction as a control.

Figure 6C shows a fluoroscopic analysis of a microchip where the SDA template was passively hybridized overnight.

Figure 7 shows the Mean Fluorescence Image of the fluoroscopic analysis of Figures 6A-6C.

Figure 8 shows a fluoroscopic analysis of a microchip where the SDA template was electronically targeted.

Figure 9 shows the titration of Factor V PCR in the SDA template of Figure 8.

Figure 10(a) shows the gel product of a NASBA amplification.

Figure 10(b) shows fluoroscopic analysis of a sandwich assay result of NASBA Tax plasmid after electronic targeting to a microarray.

Figure 11 shows a graph of the titration of non-cleavable SDA primers in Factor V anchored SDA.

Figure 12 is a schematic diagram of the anchored primers showing aspects of the branched primer design.

Figure 13 is a schematic diagram showing the stepwise process of creating amplicons from target nucleic acid sequence at a branched primer pair site.

Figure 14 is a schematic diagram showing the nature of using a signal primer to generate asymmetric ratios of nucleic acid amplicon chains such that the amplicons with signal may be electronically addressed to a capture pad for signal detection.

Figure 15 is a schematic diagram showing anchored non-branched SDA target primers.

Figure 16 is a diagram showing the layout of a microchip pad with the locations on the pad to which the various target species tested have been addressed as explained in Example 7.

Figure 17 is a photographic image of a control SDA reaction wherein no target nucleic acid was present.

Figure 18 is a photographic image showing specific localization of SDA amplified Factor V target in the presence of multiple target species on only SDA capture primer pairs specific for Factor V which had been previously addressed to only the four capture sites.

Figure 19 is a photographic image showing specific localization of SDA amplified Factor V and *Chlamydia* targets which were amplified in the presence of multiple target species and SDA capture primer pairs specific for Factor V and *Chlamydia* that had been previously addressed to specific capture sites.

Figure 20 is a photographic image showing specific localization of SDA amplified Factor V, *Chlamydia*, and Hemachromatosis gene targets which were amplified in the presence of multiple target species and SDA capture primer pairs specific for Factor V, *Chlamydia*, and Hemachromatosis that had been previously addressed to specific capture sites.

Figure 21 is a PAGE gel showing results of a multiplex solution based SDA reaction for Factor V, *Chlamydia*, and Hemachromatosis gene targets. The minus lane indicates no template DNA present, while the plus lane indicates addition of template DNA.

Figure 22 is a diagram showing a proposed reaction sequence for synthesis of a branched SDA primer pair.

Figure 23(a-c) illustrate a reaction pathway for the ligation-dependent amplification of a target nucleic acid sequence.

Figure 23(d) illustrates the ligation probes and amplification primers that would be used to detect the *Salmonella* spaQ gene present in a sample using the method illustrated in Figure 23(a-c).

Figure 24 is a graph showing specific amplification using the exonuclease ligation dependent SDA aspect of this invention, as explained in Example 10, in conjunction with a microelectrode array having capture probes for five bacterial genes pre-arranged at discrete locations.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention relates broadly to devices, methods, and compositions of matter for amplifying nucleic acid sequences in a sample and for analyzing those sequences. The amplification and the analysis are optimally accomplished using SDA and bioelectronic microchip technologies.

#### EXAMPLE 1

In a preferred embodiment of this invention, a microchip device comprising an electronically controlled microelectrode array is provided for the analysis of target nucleic acids of interest. In contrast to the uniform hybridization reaction environment and passive hybridization used in other microchip devices, the electronic microchip-based devices of the present invention offer the ability to actively transport and hybridize target and/or primer nucleic acids to capture probes at discrete locations on the surface of the microelectrode array.

Referring now to Figures 1A and 1B, a simplified version of the electronically addressable microchip-based hybridization system embodied within this invention is illustrated. Generally, a substrate 10 supports a matrix or array of electronically addressable micro-locations 12 which may be any geometric shape such as square or circular. For ease of explanation, the various micro-locations in Fig. 1A have been labeled 12A, 12B, 12C and 12D. A permeation layer 14 is disposed above the electrodes 12 and may extend over the entire device surface. The permeation layer 14 permits transport of relatively small charged entities through it, but limits the mobility of large charged entities, such as nucleic acids, to keep the large charged entities from easily directly contacting the electrodes 12 that are located under the permeation layer of a capture site. The permeation layer 14 also reduces the electrochemical degradation that could occur if direct contact were made with the electrodes 12. Electrochemical degradation is sometimes induced by both formation of reactive radical species and extreme pH at the electrode surface during the electrolytic reaction. The permeation layer further serves to minimize the strong, non-specific adsorption of nucleic acids to electrode surfaces. Attachment regions or capture sites 16 are disposed upon the permeation layer 14 and provide for specific binding sites for target materials. The capture sites 16 in Fig. 1A have been labeled 16A, 16B, 16C and 16D to correspond with the identification of the electrodes 12A-D, respectively.

The central area of the microchip contains reservoir 18 for placing sample nucleic acids above the area containing the multiplicity of capture sites 16. In a preferred embodiment, charged molecules 20, such as charged target or probe nucleic acids located within reservoir 18 may be transported to any of the specific micro-locations 12. When activated, a micro-location 12 generates the free field electrophoretic transport of any charged molecule 20 (e.g., probe, target nucleic acids or amplicons) toward the electrode 12A. As a further example, addressing electrode 12A with a positive bias and electrode 12D with a negative bias, causes electrophoretic lines of force 22 to run between electrodes 12A and 12D and further cause the transport of charged molecules 20 having a net negative charge toward the positive electrode 12A. Charged materials 20 having a net positive charge move under the electrophoretic force toward the negatively charged electrode 12D. When the net negatively charged molecules 20 contact the capture sites 16A permeation layer as a result of its movement under the electrophoretic force, the

5 charged molecule 20 becomes attached to the capture sites attachment layer 16A.  
Attachment may be by many methods as discussed below including attachment by  
10 hybridization of a target charged molecule 20 to a complementary nucleic acid probe that  
is anchored to the capture site 16.

5 Electronically addressable microchip arrays of the present invention overcome the  
size limitations of capture probe oligonucleotides and complexity requirements of passive  
15 microchip devices. The addressable microchip also greatly reduces the need for strand  
separation, at least in part, because of the use in the current system of a low ionic  
environment which inhibits the formation of double stranded nucleic acid that is in  
20 solution prior to capture and amplification of the nucleic acid at a capture site. In addition,  
the microchip arrays of the present invention allow multiple independent sample analyses  
(i.e., multiplex sample analysis) upon the same open microarray surface by selectively and  
independently targeting different nucleic acid samples to various microelectrode locations.  
25 In other words, they allow parallel multiple sample processing on an open array. As is  
described in detail below, the capability of electronic targeting to overcome the above-  
described limitations of passive hybridization methods is demonstrated in the following  
two examples A and B.  
30

#### Example A - Parallel Analysis of Single Target Nucleic Acids in a Sample

20 In a first example, a parallel analysis of the capture and detection of a single  
nucleic acid in a test sample was performed using a common locus (16S rRNA) shared by  
35 different bacterial species. Multiple comparative analyses of individual samples were used  
to identify different bacteria types.

40 The secondary structural requirements of the 16S ribosomal RNA subunit demands  
25 highly conserved nucleic acid sequences in the 16S rRNA gene. Thus, there is limited  
sequence divergence in this gene between different species of bacteria. Despite the overall  
high sequence conservation, there are pockets of microheterogeneities within the 16S  
45 rRNA gene, which can be exploited to discriminate between closely related bacterial  
species. *See, e.g., C. Woese, 51 Microbiol. Revs. 221-271 (1987).*

30 The bracketing of these microheterogeneities by conserved sequences provides  
opportunities to design many primers for consensus amplification (i.e. uniform  
50



5 amplification using the same primers regardless of species) for almost all bacterial species  
containing the conserved sequences. As shown in Figure 2A, SDA primers were designed  
10 in the conserved regions that flank the polymorphic region and used in SDA reactions.  
The resulting amplicons included the various sequences of the "microheterogeneity  
5 domains" of the 16S rRNA genes. These were analyzed by a variety of methods.

As demonstrated below, consensus SDA primers can be used for the generation of  
15 species-specific amplicons which in turn can be readily analyzed by hybridization on  
active microelectronic arrays. Similar studies have been reported using PCR as a means of  
target amplification. See, e.g., D. Linton, *et al.*, 35 *J. Clin. Microbiol.* 2568-72 (1997), M.  
20 Hughes, *et al.*, 35 *J. Clin. Microbiol.* 2464-71 (1997). However, the present invention  
uses a sandwich assay in which a single-stranded capture probe is electronically deposited  
on the array, and serves to capture one strand of a charged molecule such as a target  
25 nucleic acid or amplicon thereof. In a preferred embodiment, a multiplicity of molecules  
such as nucleic acid capture probes can be electronically deposited on different pads of the  
array. Following capture of the charged molecule to the capture sites, the captured  
15 molecule may be detected by a labeled reporter probe that binds to the captured molecule.

As is shown schematically in Figure 2A, the 16S rRNA gene near its 3' end has an  
30 oligonucleotide region stretching greater than twenty contiguous nucleotides of  
polymorphic sequence 24 flanked on both sides by conserved sequences 26. The unique  
sequences 24 of each bacterial species specified in the sequence listing herein were used in  
35 an SDA reaction in the electronically addressable microchip to show that it is possible to  
discriminate between different bacterial species by capturing these polymorphic sequences  
and their respective amplicons at specific capture sites. More particularly, primers were  
designed having nucleic acid sequence complementary to the highly conserved loop III  
40 structure of the small subunit of the bacterial ribosomal RNA 26. A 3' base  
25 complementary to a species-specific allele or point mutation in the sequence were also  
designed and made. As shown in Fig. 2A, this primer configuration facilitates design of  
both SDA amplifier and bumper primers for any particular group of organisms having the  
45 same conserved nucleic acid sequences. Primers can also be made so that they are  
30 "universal" for use in SDA to detect organisms of a group.

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In a specific example, genomic DNA from bacteria (*E. coli* O157:H7, *Salmonella typhimurium*, *Shigella dysenteriae*, and/or *Campylobacter jejuni*) were amplified. The same set of 16S rRNA encoding "consensus" primers (described in more detail below) were employed in each SDA reaction. The products of the SDA reactions were resolved on a 2% agarose gel to compare the amplification efficiencies between different bacterial species. The resulting gel is shown in Figure 2B wherein similar levels of amplification efficiency were obtained for each of *E. coli* O157:H7, *Salmonella typhimurium*, and *Shigella dysenteriae*, and in other experiments utilizing genomic DNA from *Campylobacter jejuni* (data not shown). Table I, below, shows the oligonucleotide sequences used for amplification and microarray analysis of these bacterial species.

Two different approaches were used to analyze the amplification products. A first analysis approach used a common or universal capture probe and a sequence specific reporter (*i.e.* a universal capture/specific reporter method). A second analysis approach used discriminating capture primers and a universal reporter (*i.e.* a specific capture/universal reporter method). As is shown in Figures 2C and 2D, universal capture probes 28 and universal reporters 32 were designed to span at least a portion of one of the conserved regions 26 (Fig. 2A) of the gene. As is also shown in Figures 2C and 2D, sequence specific capture probes 35 and sequence specific reporters 34 were designed to span at least a portion of the polymorphic region 24 (Fig. 2A).

Where universal capture probe 28 was used to capture nucleic acids, the initial step of hybridization between a target nucleic acid and a universal capture probe was performed electronically for several reasons. First, electronic hybridization greatly accelerates the kinetics of hybridization which is important when working with low concentrations of material, such as a highly diluted target or amplicon. Second, because of the extremely low ionic strength of the buffer systems used, targets and amplicons remain single stranded facilitating capture by probes and much less competition from the complementary strand of target or amplicon and, hence, higher net specific binding of the nucleic acid to the capture probe. Consequently, electronic hybridization allows a much higher level of nucleic acids hybridizing at the site of the capture probe resulting in greater detection and discrimination sensitivity.

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In each case of this example, reporter hybridization was passive, *i.e.* performed at elevated salt and temperature without the aid of electronics, although electronics could be used. In this particular example, since the concentration of the single stranded labeled oligonucleotides was so high, there was little practical kinetic advantage to be obtained through the use of electronic hybridization conditions. However, under different circumstances, the use of electronics during reporter hybridization may be beneficial.

As shown in Figure 3A, amplicons were addressed to the capture sites on the microchip and detected by a fluorescent reporter molecule (as described below). The relative fluorescence on capture sites to which were hybridized amplification products of bacterial 16S rRNA targets discussed in Fig. 2A were highly discriminated (*i.e.*, a polymorphism specific *Salmonella* reporter, a polymorphism *Shigella* reporter, and a polymorphism *Campylobacter* reporter). In these experiments, universal capture probes ("S") were first addressed to the microchip along with a non-specific capture probe ("NS") as a control. Amplicons from each strain-specific SDA reaction were then addressed to each corresponding row and passively hybridized with a specific reporter probe. Fig. 3A shows results for *Salmonella*-specific reporter. As a control for non-specific binding of the reporter probe to the permeation layer, a minus capture/minus target control was also performed (-C/-T). As shown, only the *Salmonella* amplicon addressed capture sites gave a positive signal. As shown in Figure 3B, not only were high discrimination ratios obtained for *Salmonella* as shown in Fig. 3A, high discrimination ratios were also seen between the various other bacterial targets. (fluorescent imaging data not shown.)

Where sequence specific capture probes 35 were used to capture nucleic acids, the initial step of hybridization between target and capture probe was also performed *electronically*. As in the universal capture example above, The reporter sequence was designed to recognize a conserved region of the 16S rDNA amplicons 26. As shown in Fig. 3C, this approach provided even higher discrimination ratios between the match and the mismatch.

#### Example B - Simultaneous Analysis of Multiple Target Nucleic Acids

In a second example, multiplex amplicon analysis was performed on the electronic microarray of the present invention. In this example, target nucleic acids from multiple

patient samples were sequentially addressed to capture sites in order to detect the presence of the human Factor V Leiden (R506Q) gene (which indicates a predisposition to activated protein C resistance and venous thrombosis). In this example, capture probes were designed so as to be specific for alleles of the R506Q gene thereby providing a method to detect allele-specific SDA.

As explained herein, since each capture site on the open microarray may be individually electronically controlled, multiple samples may be analyzed. Following amplification and position-specific targeting of each sample amplification reaction, the array was evaluated in a site-specific fashion for the presence or absence of targeted amplicons. The test system examined the presence or absence of the human Factor V Leiden mutation in several blood samples. *See, X. Liu, et al., 4 Mol. Pathol.* 191-197 (1995). The Leiden mutation is a single point mutation at the protein C cleavage site of the Factor V gene. Where this mutation has a homozygous presence in a patient, it leads to activated protein C resistance and a predisposition to deep venous thrombosis. *See, e.g., R. Bertina, et al., 369 Nature* 64-67 (1994).

To aid in discrimination, an allele-specific SDA assay was developed. The allele-specific SDA was designed to selectively amplify either the normal or the mutant Factor V Leiden allele. The SDA amplifying primers in the antisense orientation were designed with their 3' termini complementary to either the normal nucleotide base G, or the Leiden point mutation nucleotide base A, present in the sense strand of exon 10. Table I, below, shows the oligonucleotides used for amplification and microarray analysis of the Factor V gene. The corresponding sense primer was common in all reactions. However, the sense primer was modified by incorporating a biotin moiety on its 5' end in order to provide a facile mechanism for capturing any amplicons on the array following electronic targeting.

Table I: Oligonucleotides Used for Amplification and Microarray Analysis		
Bacterial 16S	Sequence (5'-3') <sup>1</sup>	Position <sup>2</sup>
BBs	CAAATGAATTGACGGGGGCC (SEQ ID NO. 1)	927-946
Bba	AAGGGTTGCGCTCGT (SEQ ID NO. 2)	1134-1120
Bas	ACCGCATCGAATGCATGTCCTCGGGTGCA TGTGGTTTAAT (SEQ ID NO. 3)	961-975

Baa	ACGATTCAGCTCCAGACTTCTCGGGTAAC <u>ATTTCACAACAC</u> (SEQ ID NO. 4)	1114-1090
Br ecoli	btr-CTCATCTCTGAAAAC TTC (SEQ. ID. NO. 11)	
Brsdys	btr-CGTATCTCTACAAGG TTC (SEQ. ID.NO.12)	
Brstyp	btr-TCCATCTCTGGATTCTTC (SEQ. ID. NO.13)	
Breje	btr-CATATCTCTATAAGG TTC (SEQ. ID. NO. 14)	
Human Factor V	Sequence (5'-3') <sup>1</sup>	Position <sup>2</sup>
FVBs	ACTACAGTGACGTGGACATC (SEQ ID. NO. 5)	
FVBa	TGTTATCACACTGGTGCTAA <sup>3</sup> (SEQ ID NO. 6)	
FVAs	bio- ACCGCATCGAATGCATGTCCTCGGGTCTC <u>TGGGCTAATAGGA</u> (SEQ ID NO. 7)	
FVA wt	ACGATTCAGCTCCAGACTTCTCGGGTAAT <u>ACCTGTATTCCTC</u> (SEQ ID NO. 8)	
FVA m	ACGATTCAGCTCCAGACTTCTCGGGTAAT <u>ACCTGTATTCCTT</u> (SEQ ID NO. 9)	
FVR	btr-CTGTATTCCTCGCCTGTC (SEQ.ID. NO. 10)	

<sup>1</sup>On amplifying primers, BsoBI recognition sites are boldfaced, genomic homology regions are underlined and Factor V allele-specific 3' termini are shown in italicized boldfaced type. The designation "bio" represents biotin conjugation and "btr" indicates fluorescent BODIPY Texas Red conjugation. FVR is reporter for Factor V, FVAs is sense strand for amplification while FVAwt and FVA<sub>m</sub> are amplification primers for wildtype and mutant respectively. FVBs and FVBa are sense and antisense bumper primers for Factor V.

<sup>2</sup>Bacterial 16S sequence was obtained from GenBank, human Factor V sequence refers to GenBank accession #L32764.

In this multiplex Factor V gene study, four clinical DNA samples were analyzed in duplicate without prior knowledge of the patient's Factor V Leiden mutation status. Two allele-specific SDA reactions were conducted per sample (containing either normal or mutant primers) to examine each patient's genotype. The amplifications were conducted in parallel. The PAGE results from five of these pair-wise reactions is shown in Figure 4A wherein the allele-specific amplification reactions under these conditions are shown to be highly specific. That is, the selective absence of visible mutant or normal-type amplicons indicates that the amplification reaction is sensitive to the presence or absence of the Factor V Leiden mutation in these individuals.

All amplicon reactions, regardless of the presence or absence of amplified material as determined by gel analysis, were uniformly treated and sequentially targeted to specific locations upon the microarray. Representative results from three DNA patient samples are shown in Figure 4B. These samples were targeted in duplicate. The presence or absence of a fluorescent signal from a hybridized reporter oligonucleotide complementary to a conserved region on the target amplicon (*i.e.*, a "universal" reporter probe) indicates the presence or absence of Factor V amplicons. As can be seen, the fluorescent signal correlates well with the gel results shown in Figure 4A.

As is shown in Figure 4A, positive signals were several fold greater than background signals. In general, the true mutant signal was lower than that from wild type amplicons (as shown in Figure 4B). The sites were scored simply by making the criteria for a positive signal to be at least twofold above the background fluorescence present at non-addressed capture sites. As shown in Table II, below, there was complete correlation between the presence of amplified material by gel analysis and the presence of strong or moderate fluorescent signals upon the array.

The strength of the fluorescent signal approximated the *apparent quantity* of amplified material. This was most striking in those samples with an apparently less efficient amplification reaction such as was seen with the DNA from patient 961a in Figures 4A and 4B. In short, these results show that multiple sample analysis by the serial application of samples followed by single reporter detection works using a microelectronic

array, and shows that this process may serve to supplement or replace other forms of analysis, *e.g.* gel electrophoresis, in the same or similar analyses.

Since these samples were analyzed prior to knowledge of their mutational status, it was of interest to determine whether the apparent allele specificity of the amplification reaction did, in fact, correspond with clinical status. As is shown in Table II, below, the selectivity of the allele-specific amplification reaction was in complete agreement with the Factor V Leiden mutational status of each sample as determined by PCR and MnlI restriction site analysis. (R. Press, unpublished observations). Thus, combined with allele-specific SDA, analysis of amplicon product formation upon an electronically addressable array is a useful method for detecting genetic point mutations in multiple patient samples.

Table II: Allele Specific Factor V SDA Amplification Results

Patient	Sample Date	PAGE <sup>1</sup>		Microarray <sup>1</sup>		Genotype <sup>2</sup>
		wt	mut	wt	mut	
951961	04/10/97	X	X	X	X	Heterozygous
951961	06/04/97	X	X	X	X	Heterozygous
952018	04/10/97	X	O	X	O	Homozygous wt
952018	06/04/97	X	O	X	O	Homozygous wt
960286	04/10/97	O	X	O	X	Homozygous mut
960286	06/04/97	O	X	O	X	Homozygous mut

<sup>1</sup>"X" indicates positive, "O" indicates negative.

<sup>2</sup>Genotype was determined by PCR-RFLP with Mln-I restriction enzyme by methods well known to those skilled in the art.

Experimental Protocol used in the above described data

**Materials** – Deoxynucleoside 5'-triphosphates (dGTP, dATP, TTP) were purchased from Pharmacia, Alameda, California. 2'-deoxycytosine 5'-O-(1-thiophosphate) (dCTP $\alpha$ S), BsoB1 restriction endonuclease and Bst polymerase were supplied by Becton Dickinson, Sparks, Maryland. Oligonucleotides were synthesized by Oligos, Etc., Wilsonville, Oregon.

5  
10  
5  
15  
20  
10  
20  
SDA Amplification – Amplification reactions utilized either 1 µg of genomic DNA (16S) or 0.1 µg of genomic DNA (Factor V) in a volume of 30 µl. Amplification conditions and concentrations were adapted from that presented previously (see, C. Spargo, *et al.*, 10 *Molecular and Cellular Probes* 247-256 (1996)) with the following changes: The 5'to3' exonuclease deficient polymerase Bst replaced the exo-BCA polymerase, as disclosed and used in M. A. Milla *et al.*, *Biotechniques*, v24, p 392-396, March 1998 herein incorporated by reference. For 16S amplification, 25U/reaction (Bst) and 60U/reaction (BsoB1) were used. Oligonucleotides employed for amplification reactions are shown in Table I above. Reactions were allowed to proceed for 30 minutes at 60°C and then terminated by the addition of 10 µL of 100 mM EDTA and then stored at -20°C.

25  
15  
Gel Electrophoresis – Amplification reactions were analyzed using standard protocols with either 1% agarose gel or with 6% polyacrylamide mini gels (Novex, San Diego, California) followed by ethidium bromide staining. Images were obtained using an AlphaNotech Chemimager (San Leandro, California).

30  
20  
35  
40  
25  
45  
30  
50  
55  
Electronic Microarray Analysis – The microelectronic array assembly has been described previously. See, R. Sosnowski, *et al.*, 94 *J. Proc. Natl. Acad. Sci. USA* 119-123 (1997). Electronic targeting of capture oligonucleotides (biotin-GGATGTCAAGACCAGGTAAGGTTCTTC, Genbank locus 988-1014 bp (SEQ ID NO. 15) and hybridization of amplicons (16S) or reporter oligonucleotide (Factor V) utilized conditions reported elsewhere. See, R. Sosnowski, *supra*, and C. Edman, *et al.*, 25 *J. Nucleic Acids Res.* 4907-4914 (1997). In brief, crude amplification reactions were either spun for two minutes through G6 columns (Biorad, Hercules, California) preequilibrated with distilled water or dialyzed in multiwell plates (Millipore, Bedford, Massachusetts) for more than or about five hours against distilled water. The prepared samples were then mixed in a 1:1 ratio with 100mM histidine and heated at 95°C for five minutes prior to electronic addressing. For analysis of 16S amplicons, electronic hybridization of the amplicons was performed, followed by hybridization in 6X SSC of a fluorescent labeled oligonucleotide reporter homologous to a specific bacterial sequence. Specific nucleotide sequences are shown in Table I, above. Passive hybridization was allowed to proceed for 30 minutes at room temperature. The microchips were washed 5 to 8 times using 0.1X



STE/1% SDS followed by 1X STE. Similar conditions were employed for the single target experiment above using the 16S bacterial rRNA sequence-specific Biotin-captures and a common btr-labeled reporter for detection. For analysis of Factor V amplicons, a fluorescent-labeled oligonucleotide (btr-CTGTATTCTCGCCTGTC (SEQ ID NO. 10) was introduced in 6X SSC and allowed to hybridize for 30 minutes at room temperature. The array was then washed in 0.1X STE/1% SDS followed by 1X STE.

#### EXAMPLE 2

Turning now to the electronic amplification aspect of the present invention, target nucleic acid is electronically concentrated in the vicinity of anchored primers located on a capture site and used in an SDA or other amplification method. The target nucleic acid may be electronically concentrated and hybridized to binding molecules (e.g., capture probes) on the surface of the microchip capture sites prior to the introduction of SDA reaction components (i.e. enzymes, nucleotides, etc.) thereby increasing the efficiency and decreasing the time necessary for hybridization of target nucleic acid to the anchored capture primer on the capture site. Hybridizing the target nucleic acid to specific locations on the microarray prior to addition of SDA reaction components also permits the array surface to be washed to remove unwanted and possibly interfering non-target nucleic acids from the reaction environment. Thus, amplification reactions, such as anchored SDA, can benefit greatly by using an electronically addressable microarray system.

The components of the amplification reaction itself (without template and amplification primers) are introduced and the amplification reaction allowed to proceed. There are at least three advantages to employing electronic targeting of template molecules. The first is that the overall time and efficiency of the amplification process is dramatically improved since a major rate-limiting step (that of the time required for the template to find the anchored primers) is removed from the overall reaction rate. Also, the use of the electronic concentration and hybridization increases the number of target molecules at the selected site, as compared to non-electronic passive hybridization for an equivalent time period, thereby increasing the *absolute* numbers of starting template molecules for amplification resulting in improvement in both the overall yield of the amplification process and the sensitivity of the system to lower starting template numbers.

5  
10  
15  
The second advantage is that discrete target nucleic acid samples can be applied to specific locations upon the array surface thereby allowing multiple and different nucleic acids to be amplified simultaneously on one array. Alternatively, a nucleic acid may be targeted to several different locations, each containing specific sets of amplification  
5 primers so that multiple different amplification reactions can be simultaneously carried out from a single sample. As noted above, the ability to remove unnecessary or unhybridized nucleic acids from the reaction mixture significantly aids this process.

20  
25  
A third advantage to this approach is that following the amplification reaction, the captured amplicons are available in a site-specific fashion for subsequent analyses, either  
10 by introduction of fluorescently labeled nucleotides or by the incorporation of labeled oligonucleotides during the course of the amplification reaction or by hybridization with an appropriate reporter oligonucleotide at the end of the reaction by denaturation of the amplicons that are bound to the capture sites.

30  
35  
40  
In an example of this electronic addressing embodiment, an experimental protocol was designed to *enhance* anchored Factor V SDA sensitivity by using electronic hybridization of Factor V encoding template nucleic acid to anchored SDA primers (Seq. I.D. Nos. 20 and 21) on a microchip array. The SDA primers were biotinylated at their respective 5' ends. These primers also contained a BsoBI enzyme cleavage site. The reaction mix included the bumper primers (Seq. I.D. Nos. 22 and 23) for SDA. The  
20 microchip array was prepared by scraping the streptavidin-agarose layer from the outer electrodes of the microchip. The edges of the chip were waterproofed with Rain-X and the surface was buffed clean with a cotton swab applicator. The array was incubated with milli-Q water for at least 30 minutes at room temperature.

45  
50  
55  
Solutions were prepared for electronic addressing on the microchip. SDA primers  
25 in 1  $\mu$ M in 50 mM histidine buffer, 1  $\mu$ M biotinylated T12-btr oligonucleotide in 50 mM histidine buffer, and 50 mM histidine wash buffer were prepared. The microchips were washed with 50 mM histidine buffer, and biotinylated T12-btr oligonucleotides were addressed using a standard A/C protocol (800 nAmps for 25 seconds) to selected capture sites to check the quality of the streptavidin microchips. The SDA primers were addressed  
30 to selected capture sites as shown using the standard A/C protocol.

5  
10  
5 For electronic hybridization (as opposed to passive hybridization) experiments, double stranded PCR nucleic acid templates were first denatured at 95 °C, and an equal volume of 100 mM histidine buffer was added to the template. The template mixture was then electronically hybridized to the capture SDA primers using a standard A/C protocol for hybridization (1.6 µAmps, 60 seconds).

15  
10 For passive hybridization experiments, asymmetric PCR nucleic acid templates were first denatured at 95 °C for 5 minutes. The solution was then brought to a 4X SSC concentration with a 20X SSC (3M NaCl, 0.3 M NaCitrate) stock and 20 µl of the mixture was pipetted onto a microchip (which had been previously electronically addressed with SDA primers) and incubated at room temperature overnight.

20  
25 After incubation the microchip arrays were washed 2X with water and incubated with 1 mg/ml BSA for 30 minutes at room temperature to block any non-specific binding sites. The microchips were washed again with water (2X) and pre-warmed at 60 °C for 5 minutes. All SDA solutions were also pre-warmed at 60 °C for 5 minutes. After pre-warming, the water was removed from the microchips and incubated with 10 µl SDA reaction mix (40 mM K<sub>2</sub>HPO<sub>4</sub> pH 7.6, 1.6 mM each dCTP<sub>α</sub>S, dTTP, dATP and dGTP, 8.3 mM MgCl<sub>2</sub>, 1.3 units BsoBI and 0.5 units Bst polymerase) for 30 minutes at 60 °C in a humidifying chamber. The reaction was stopped by removing the supernatant from the microchip surface to an eppendorf tube containing 2 µl of 100 mM EDTA.

30  
35  
40  
25 After the SDA reaction, the microchips were washed 3X with 0.5X SSC, pH 7.2. The SDA products were then denatured on the microchip *in situ* with addition of 0.5 X SSC, pH 12.0 for 4 minutes, washing the microchip with additional buffer after every minute. The microchips were then washed with 0.5X SSC, pH 7.2 at least 3 times, then with 4X SSC, pH 7.2 at least three times. The microchips were incubated with a 1 µM mix of btr-labeled reporter oligonucleotides (such as Seq. I.D. Nos. 24 or 44) in 4X SSC for 3 minutes at room temperature, washed extensively with 4X SSC at room temperature, then imaged.

45  
50  
30 For passive hybridization of Factor V template on microchips addressed with Factor V SDA primers at distinct sites, microchips were addressed with 1 µM of either Factor V SDA primers, or Factor V SDA primers lacking a BsoBI site as a negative control for the SDA reaction. Since the negative control lacks a BsoBI site, the reaction

can only undergo primer extension upon binding of a template and not SDA amplification.

This reaction controls for the presence of non-specific binding as well as the production of non-specific amplification products with which the reporter oligonucleotides may react. A no-template control was also present. These microchips were then fluoroscopically analyzed for Factor V amplicons having the fluoroscopically labeled btr-reporter oligonucleotides. SDA products were seen only in the microchip where SDA template was *passively* hybridized *overnight* (Fig. 6C). No products were seen in the no-template control microchip (Fig. 6A), or in the microchip (Fig. 6B) where BsoBI was not included into the reaction (another negative control for the SDA reaction). In the microchip that was passively hybridized (Fig. 6C), the SDA products are seen only in the area where the SDA primers were addressed, not in the non-cleavable SDA primer quadrant of the array, again confirming that the product detected is specific and is driven by an SDA-based process. The drawback of this assay is that the images seen after the SDA reaction were very weak, having MFI (Mean Fluorescence Image) values of 14 at an integration time of 1s for non-diluted template levels (Fig. 7).

For *electronic* hybridization of Factor V template to anchored SDA primers on a microchip, experiments were conducted in a manner parallel to that carried out for passive hybridization, with the exception that hybridization of the template was facilitated by electronic addressing. Additionally, the template was also serially diluted. As a control, passive hybridization of the factor V template was carried out and resulted in a very small increase (approximately 1 MFI unit) over background in the SDA reaction. Again, no signal was seen in the non-cleavable primer quadrant, indicating the need for SDA-directed amplification in this system. In contrast, the microchip that was electronically hybridized showed a signal in the SDA primer quadrant (Fig. 8) and showed a significant signal in all dilutions tested (Fig. 9). Even at a dilution of 1:100 of the Factor V template, the signal was still very high, at approximately 19.4 MFI/sec. Given that the MFI signal of a 1:100 dilution of the electronically addressable microchip was 19.4 times higher than the signal from a passively hybridized chip in this experiment, (and 1.4 times higher than in the passive hybridization experiment, above, where the template was not diluted) the efficiency of the SDA assay increased approximately 140-1940 percent by using an electronic hybridization protocol. This demonstrates that electronic hybridization of the

5  
10  
template to SDA primers anchored on the microchip increases the sensitivity of the assay approximately 1000 fold. In addition, the time required to perform the entire SDA experiment was reduced by one full working day (as compared to passive hybridization wherein the template needed to be incubated overnight to achieve efficient binding levels).

5  
15  
In another example, we show that electronic addressing of target molecules to capture sites facilitates the amplification of DNA or RNA target nucleic acids using the technique known as nucleic acid sequence-based amplification (NASBA). In this method three different enzymatic activities are used in a coordinated fashion with an isothermal method of amplification. In this electronically-mediated process, the simultaneous or  
20  
10  
multiplex amplification of different sequences is possible either by site specific targeting and amplification or by using multiple primer sets. Moreover, NASBA, as practiced in the invention, may use either anchored or solution-based primers in the amplification reaction. In either case, the reaction is enhanced using electronic addressing of the target to its  
25  
respective amplification primers.

15  
30  
In this example, target nucleic acid sequences were first electronically hybridized to discrete locations upon a microchip. Unwanted or non-specifically binding nucleic acids were removed either by electronic washing, or passive (non-electronic) washing or by a combination of the two. Following the wash step, the hybridization solution was replaced by a buffer cocktail comprising amplification primers, nucleotides, magnesium chloride and the enzymes or enzymatic activities necessary for amplification. (These  
20  
35  
enzymatic activities are: reverse transcriptase activity; RNase H activity; and RNA polymerase activity. The activities of these enzymes coordinately serve to amplify the isolated sequences in a fashion similar to that of NASBA.) Once the amplification cycles were completed, the amplified material was electronically isolated or captured and then  
40  
25  
quantitated (*i.e.*, detected) by various methods known in the art. In general, such detection may be carried out using, for example, a capture oligonucleotide specific for the newly synthesized region or, a fluorescently-labeled oligonucleotide in a "sandwich assay."

45  
50  
Each stage of this process is augmented as compared to existing technology. For instance, the electronic targeting of the target sequence followed by its specific  
30  
hybridization using suitable capture oligos (*e.g.* the primers for amplification) allows for the electronic removal of unwanted or contaminating DNA or RNA. The removal of

5 nonspecific nucleotides that can cause non-specific binding and amplification, allows for a  
higher complexity of amplification events to simultaneously occur, as well as for more  
10 specific amplification. In addition, if all the primers for amplification are anchored,  
amplification events using different target sequences can occur simultaneously at different  
5 locations upon the chip or device, *i.e.* multiplex reactions. The enzymes themselves can  
also be targeted, allowing for greater precision in mediating the amplification events or  
15 stages. Finally, the products of the amplification reaction can also be targeted to  
alternative sites and quantified, allowing the progress of the amplification reaction to be  
followed.

20 10 In one representative, but not limiting experiment, NASBA amplification of an  
HTLV1 plasmid was performed in solution using three different concentrations of  
template plasmid (approximately 1 ng, 1 pg, and 1 fg). The reaction employed an initial  
melting of the DNA template at 95 °C, followed by an isothermal annealing step of 15  
25 minutes at 50 °C. The annealing reaction consisted of 8 µl of 2.5X NASBA mix (100 µL of  
15 25 mM NTP mix, Pharmacia Lot #60920250111; 50 µL of 25 mM dNTP mix, Pharmacia  
Lot #6092035011; 50 µL of 1M Tris, pH 8.5; 31.25 µL of 2M KCl; 15 µL of 1M MgCl<sub>2</sub>;  
and 253.75 µL sterile H<sub>2</sub>O), 1 µl of a 5 µM concentration of an oligonucleotide primer  
30 (#885; 5' AATTCTAA TACGACTCAC TATAGGGAGA GGTGATCTGA  
TGTCTGGAC AGG 3' (SEQ ID NO. 16), and 1 µl of one of the three dilutions of the  
20 HTLV1 plasmid (or none) in four separate tubes to achieve 1 ng, 1 pg, 1 fg, and 0 final  
concentrations. Enzymes which would not survive the 95 °C denaturation step were added  
35 at the beginning of the amplification step. Thus, 1 µL of 100 mM DTT (dithiothreitol) and  
then 0.5 µL AMVRT (AMV reverse transcriptase from Boehringer Mannheim (Cat No.  
1495 062; Lot No. 83724624-76) were added at the 50 °C step. The reaction was  
40 terminated by heating to 95 °C for 5 minutes. The tubes were placed on ice.

25 Following the annealing reaction, an amplification reaction was set up, also in four  
tubes, consisting of 10 µL 2.5 X NASBA mix; 1 µL of 250 mM DTT; 0.3 µL of Rnase H  
45 (Ribonuclease H from Boehringer Mannheim, Cat No. 786 349; Lot No. 13656445-05);  
2.5 µL enzyme mix (20u T7 polymerase from Boehringer Mannheim Lot # 83495822-31;  
30 8u AMV RT; 0.2u RNase H; and 2.5 µg Rnase and Dnase free BSA (Bovine Serum  
Albumin) from Pharmacia # 6078914011); 6 µL of primer mix (5 µL of 5 µM primer  
50

5 #885; 5  $\mu$ L of 5  $\mu$ M primer #882: ACTTCCCAGGGTTTGGACAGAGT (SEQ ID NO.  
17); 18.75  $\mu$ L 100% DMSO; and 1.25  $\mu$ L H<sub>2</sub>O); and 2  $\mu$ L of primed DNA from the four  
10 annealing reaction tubes, each placed in a separate tube. The reaction was incubated for 60  
minutes at 40 °C, then put on ice. The reactions (10  $\mu$ L) were then separated on a 2%  
5 agarose gel and stained with ethidium bromide.

15 The highest concentration of starting template plasmid produced the largest amount  
of product, whereas the lower two concentrations produced little or no product (Fig. 10a).  
The product of the 1 ng reaction (the bright band on the gel) was cut out of the gel and  
then diluted either 200-fold, 500-fold or 1000-fold in 50 mM histidine. The reaction  
20 product of the 1 pg template reaction was also diluted 200-fold for comparison. These  
reaction product dilutions were then electronically targeted to capture sites upon a  
microarray containing either specific (500  $\mu$ M of XL5R.bio, 5'  
25 TTCTTTTCGGATACCCAGTCTACGTGTTTG 3' (SEQ ID NO. 18) or non-specific  
(ATA5.bio) pre-targeted capture antibodies using 500  $\mu$ A constant current for 1 minute,  
15 changing the buffer and targeting the next capture site without washing. After targeting  
the reaction products, the capture sites were washed 5X with histidine (50 mM) and the  
fluorescence at each location evaluated (Fig. 10b) using a fluorescently labeled reporter  
30 oligo (HTVPXs.313TR; (NH<sub>2</sub>)-ACTTCCCAGGGTTTGGACAGAGT 3' (SEQ ID NO.  
19) 15  $\mu$ L) introduced, passively, for 15 minutes at 25 °C. Following 3 washes in 1 mL of  
20 0.2X STE/1% SDS, and 5 additional minutes in STE/SDS, the capture sites were rinsed,  
35 and a 2 second image was taken. Thereafter, the buffer was changed to histidine and the  
capture sites were run by column at 200 $\mu$ A/pad for 1 minute, washed, and a 2 second  
image taken. The results of the electronic sandwich assay of the amplified reaction  
40 paralleled the relative amounts of amplified product introduced, as shown in Figure 10b.

### 25 EXAMPLE 3

45 In yet another example, anchored SDA is carried out, preferably using electronic  
targeting of the target nucleic acid to the specific site, and preferably including at specific  
sites upon the array *non-cleavable* oligonucleotides in combination with a greater ratio of  
30 normal SDA primers (*i.e.* the non-cleavable primers do not contain the requisite restriction  
endonuclease site necessary for SDA, but which are identical to SDA primers in other  
50

5 aspects). Anchored SDA is then carried out, using electronic targeting of template nucleic  
acid to the specific site followed by amplification and reporter hybridization. The optimal  
10 ratio of non-cleavable to normal primers is determined empirically, and is based on the  
signal obtained from reporter labels. Alternatively, other sites and/or functionalities can be  
5 introduced upon these non-amplifying primers for the purposes of subsequent cleavage and  
analysis or other manipulations. The prime criteria of these non-cleavable primers is that  
15 the 3' terminus contains sufficient homology to the target nucleic acid or amplified  
products thereof to hybridize and serve as the basis for primer extension by polymerase.

In a specific experiment of this example, different proportions of standard Factor V  
20 10 amplifying primers were mixed with primers which no longer had a BsoBI site present.  
These mixtures were targeted to different locations upon the array and diluted Factor V  
PCR amplicons were targeted to each location. The entire array was then washed and a  
25 mixture containing SDA amplification reaction components (except amplifying primers)  
was added. The amplification reaction was allowed to proceed for 30 minutes at 60°C  
15 then, following denaturation, Bodipy-Texas Red labeled reporter probes were added and  
hybridized. The fluorescence present at each site was then quantified.

30 The experimental protocol followed in this experiment was as follows. First,  
microchips were prepared for electronic addressing and hybridization by scraping any  
agarose away from the outer electrodes and treating each microchip surface with Rain-X.  
20 20 The chips were washed three times with water and allowed to stand in water for at least  
35 about 30 minutes. Then Factor V SDA primers (*i.e.*, Seq. I.D. Nos. 20 and 21) and non-  
cleavable (NC) primers (*i.e.*, Seq. I.D. Nos. 42 and 43) were diluted to 2  $\mu$ M total (from 0-  
100% non-cleavable primers mixed with SDA primers, see Table III below) and equal  
40 volumes of 100 mM histidine were added to make a 1  $\mu$ M primer solution in 50 mM  
25 histidine buffer. Next 10 nM btr-T12 and 1  $\mu$ M ATA-5 oligos were prepared as controls  
in 50 mM histidine. Factor V template DNA was then diluted to an appropriate  
45 concentration and incubated at 95.5°C for about 5 minutes. An equal volume of 100 mM  
histidine was added to make a final concentration of 50 mM histidine buffer.



**Table III: Non-Cleavable Primers to SDA Primers Mix**

% Non-Cleavable Primers	2 $\mu$ M NC Primers ( $\mu$ l)	2 $\mu$ M SDA Primers ( $\mu$ l)
0	0	100
10	10	90
20	20	80
30	30	70
40	40	60
50	50	50
60	60	40
70	70	30
80	80	20
90	90	10
100	100	0

The SDA/Non-Cleavable primers mix, as well as controls, were then electronically addressed and a template was hybridized onto each microchip array. An image was taken and the microchips were washed three times with water and incubated with 1 mg/ml BSA for 30 minutes at room temperature. The microchips were then washed two times with water and pre-incubated at 60°C for 5 minutes in a humidifying chamber (*i.e.* a petri dish with moistened Whatman 3 MM paper).

An SDA mix comprising 40 mM  $K_2HPO_4$ , 1.6 mM dCTP $\alpha$ S, 1.6 mM dTTP, 1.6 mM dCTP, and 1.6 mM dGTP, 8.3 mM  $MgCl_2$ , 1.3 units BsoBI enzyme, and 0.5 units Bst polymerase, was pre-incubated at 60°C for 5 minutes. Water was removed from the microchips and 10  $\mu$ l of pre-warmed SDA mix was added to each microchip without allowing the microchips to cool down. The microchips were then incubated at 60°C for 30 minutes. The SDA reaction was then stopped by removing the solution from each microchip and transferring it to an eppendorf tube containing 2  $\mu$ l 100 mM EDTA. The supernatant was then analyzed on non-denaturing polyacryamide gels.

The microchips were washed with 0.5X SSC solution, wherein the SSC solution comprises 75 mM NaCl and 7.5 mM NaCitrate, pH 7.2, at least three times. Next, the microchips were incubated in 0.5X SSC, pH 12 solution for 4 minutes, with the solution being pipetted up and down about every minute. Each microchip was washed at least three times with 0.5X SSC, pH 7.2, then three times again with 4X SSC solution. Passive hybridization of 1  $\mu$ M reporter oligonucleotides in 4X SSC was then carried out at room temperature for 3 minutes. Each microchip was washed extensively with 4X SSC. If necessary, an additional stringent wash with 0.2X SSC/1% SDS was done for 5 minutes at room temperature. The microchips were then washed extensively with 0.2X SSC. Finally, the microchips were imaged with appropriate lasers and filters, and the fluorescence present at each site was quantified. Results from this experiment are shown in Fig. 11. As shown in Fig. 11, a 10% optimal percentage of non-cleavable SDA primers included in the SDA primer mix for anchored SDA gave an approximately 2-fold increase in specific signal over the absence of non-cleavable primers (0%). As expected, with an increase in non-cleavable to SDA primer ratios, the efficiency of the SDA reaction decreases to levels where no detectable SDA amplification can be seen. This demonstrates that the addition of non-cleavable primers to the SDA primer mix, which in effect retains any signal that may have been nicked prior to denaturation of the double-stranded template, improves signal intensity in anchored SDA.

#### EXAMPLE 4

In another embodiment, an amplification method of the present invention comprises an allele-specific SDA method. The method preferably selectively amplifies only those strands that include a specific allele. The method preferably utilizes amplifying primers designed so that their respective 3' termini include nucleotide bases that are complementary to the nucleotide sequence of the desired allele. At least one of the primers may also preferably include a biotin moiety on its 5' end to provide a facile mechanism for capturing amplicons on the array following electronic targeting and amplification. Generally, the specificity of the process of this example is derived from the low efficiency of nucleic acid chain extension when the 3' terminal nucleotide of the primer is non-complementary to the target sequence.

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In a modification of this example, individual amplified patient nucleic acid samples are immobilized in discrete locations on the microarray, and all samples are probed simultaneously with gene or allele-specific reporter probes. Individual patient samples are immobilized by introducing biotin into the samples during SDA. One of the SDA primers is added which contains a 5' biotin linker which does not have a restriction cleavage site, and, therefore, is not cleavable. The samples are denatured and addressed to individual capture sites. A single stranded amplicon from each patient is immobilized at an individual capture site. Once all patient samples are immobilized, they are all probed simultaneously and in parallel. Thus, an open microchip is used to analyze multiple patient samples with minimal cross-contamination.

10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In this example, the biotinylated primer is preferably either a noncleavable version of the flanking primer used for amplification, or an internal sequence. In either case, it forms a dead end product (*i.e.* one which is not further amplified). The primer is preferably present in limited amounts so that the entire primer is converted to product. For instance, when screening for a genetic mutation such as, for example, the Factor V Leiden mutation, there are only two alleles, a wild type and a mutant. Amplification is performed using primers which are specific for the wild type locus, but not the allele (*i.e.*, mutant). The internal biotinylated primer is converted to a product shorter than the full length amplicon through extension if the allele is present. The fragment is then addressed to a pad and subsequently probed with an allele-specific probe, or an allele-specific biotinylated internal probe is used. Amplification may take place in the presence of fluorescently labeled nucleotides. Preferably, each patient sample is amplified in two separate reactions with allele-specific primers (for wild and mutant alleles) which are then addressed to different pads, or the two reactions are performed simultaneously using reporter molecules that fluoresce in two different colors and both products are addressed to the same capture site (in which case genotype would be determined by the fluorophore remaining at the site for that patient).

(See Example 1 B for additional embodiments of allele-specific methodology and technique).

## EXAMPLE 5

In another embodiment, SDA products may be simultaneously generated and specifically captured on a microchip by performing thermophilic SDA (tSDA) in a flow cell region over the microchip of the present invention. (see U.S. Patent No. 5,648,211 for a discussion on tSDA and U.S. Patent No. 5,547,861 for a discussion on signal primer extension, both herein incorporated by reference). By tSDA is meant SDA using thermophilic enzymes allowing operation at temperatures in excess of 40°C to facilitate stringent hybridization. Prior to amplification an internal capture sequence having a 5' biotin modification is immobilized preferably to a specific streptavidin-containing capture site location. As single stranded amplicons are generated free in solution during the SDA process, a fraction of the amplicons specifically hybridize to the immobilized capture oligonucleotide. Detection of the hybridized strand is preferably via one of the methods described throughout this disclosure. This embodiment of the method allows use of very small sample volumes (e.g., on the order of about 10  $\mu$ l), and allows for specificity controls due to use of sequences for capturing that are preferably located on separate capture sites and are internal to the sequences used to perform SDA priming. Moreover, detection of the captured sequences may occur in "real time" as they are being generated during the SDA reaction thereby facilitating the simultaneous SDA and monitoring of the SDA reaction and generated amplicons.

With respect to this method there are two exemplary schemes to incorporate a fluorescent species for detection. In a first scheme to incorporate a fluorescent species for detection, as is shown in Figure 5A, an additional oligonucleotide 36 is included in the amplification reaction. This additional oligonucleotide is fluorescently labeled and binds to its single stranded complemer generated by the amplification process. Upon binding, polymerization is initiated in a 5' to 3' direction from this primer by the polymerase 37 used in the SDA reaction. As a course of the regular amplification process, an oligonucleotide which functions as an amplifying primer binds 5' upstream to the same strand as the fluorescently labeled species. As polymerase extension occurs from this primer, the fluorescently labeled strand is displaced and released as a single stranded species free into solution above the array. On the array are previously addressed anchored complementary oligonucleotides. These serve to capture a portion of the fluorescently

5 labeled oligonucleotides and provide a fluorescent signal upon the array which is both  
location-specific (and, therefore, sequence specific) and increasing over the course of the  
10 reaction.

5 In a second scheme to incorporate a fluorescent species for detection, as is shown  
in Figure 5B, anchored capture oligonucleotides have either an unmodified endonuclease  
restriction sequence and capable of supporting an SDA reaction 40 or a modified sequence  
15 that will not be recognized by an endonuclease 45. These anchored capture primers 40 and  
45 are used to bind single stranded products 42 of the amplification reaction. These  
capture oligonucleotides 40 and 45 serve as the site for oligonucleotide extension by  
20 polymerase activity. Upon completion of the amplification reaction, the double stranded  
material is melted, preferably by electronic or chemical methods (including, for example,  
alkaline in pH 12), releasing the original amplicon 42 and extension product 43. The array  
is washed and then a fluorescently labeled oligonucleotide 44 is introduced. These  
25 reporter oligonucleotides specifically hybridize only to the polymerase extended portions  
of the capture oligonucleotides 40 and 45. In this scheme it is preferred that the ratio of  
cleavable to noncleavable oligonucleotides is about 10:1. It is believed that this ratio  
allows the amplification reaction to optimally proceed while providing a sufficient number  
30 of uncleaved extension products remaining at the capture sites for detection by reporter  
probe.

#### 35 EXAMPLE 6

In still another embodiment of the invention, SDA is preferably conducted directly  
on an electronically addressable microchip under the following conditions. The sample is  
40 initially prepared and randomly sheared to less than about 5kB. The sample is then  
denatured and target nucleic acid is captured to a single capture site that contains both 5'  
25 and 3' SDA primers. "Bumper primers" which hybridize to the regions immediately  
upstream of the capture primers are added in a relative concentration of about 1/10 that of  
the capture oligonucleotides. An SDA mix (*i.e.* 3 unmodified dNTPs, 1 thiol modified  
45 NTP, (and, possibly, a fluorescent labeled NTP,) and enzymes preferably comprising  
thermophilic exo (-) DNA polymerase plus restriction enzyme) are passively added. The  
30 microchip is then heated to about 40-60°C and SDA is allowed to proceed.

“Real time” detection of the SDA reaction and product amplicons is possible by incorporating NTPs which allow fluorescent energy exchange or quenching. For example, an NTP containing Bodipy Texas red is combined with one that contains Cy5. Incorporation of NTP via polymerase elongation can be continuously monitored by monitoring fluorescent energy shift.

Under one theory it is believed that the preferred minimum estimated spacing between adjacent oligonucleotides on a pad is about 1.25 nm ( $10^4$  ODNs/ $80 \mu\text{m}^2 = 100 \times 10^6 / 80 \times (10^3)^2 \text{ nm}^2 = 1.25 \text{ ODN/nm}^2$ ). If oligonucleotide bridging is required to start SDA, then it is believed that the optimal length of an SDA fragment which will allow optimal amplification can be determined empirically. As a starting point, 100bp = 34 nm seems reasonable.

#### EXAMPLE 7

In this embodiment, a novel method of an anchored SDA reaction which alters the spatial relationships between amplification primers, target DNA, and enzyme molecules is provided. Because both amplification primers are brought into close proximity to one another, the efficiency of the SDA reaction is actually increased. The spacing relationship between the amplification primers may also be adjustable by altering linker elements between the primers thereby enabling precise definition of the stoichiometry ratios of the primers, the local concentration of the primers, site directed template capture, and spatial relationships of the primers, so as to set up the SDA mechanism in a coupled-concerted fashion to benefit exponential amplification of target DNA.

Referring now to Figs. 12 or 15, SDA target capture primers are attached to specific areas or capture sites 5 on an electronically addressable microchip. The capture primers are attached at each site such that both upstream and downstream primer pairs required for SDA specific for a target nucleic acid of interest are present together in close proximity to one another at the capture site. With regard to Fig. 12, branched structure 3 is attached to capture site 5 and to the 5' ends of plus and minus strand SDA nucleic acid primers. For each primer, an unmodified restriction site sequence 1 (i.e., the unmodified strand of a hemimodified restriction site) is located 5' to target specific capture sequences 2 and 4. With regard to Fig. 15, linear plus and minus strand nucleic acid SDA primers are

5 attached to capture site 5 at their respective 5' ends. Like the branched primer pairs, the  
linear SDA primers comprise unmodified restriction site 6 sequence 5' to target specific  
10 capture sequences 7 and 8.

The microchip may be prepared according to teachings known in the art such as the  
5 method disclosed in US Patent no. 5,605,662 herein incorporated by reference. In the  
current example, prior to addition of SDA primers, the streptavidin-agarose layer was  
15 scraped from the outer electrodes of the microchips. The edges of each microchip were  
waterproofed with Rain-X (Unelko Corporation, Scottsdale, Arizona) and the surface of  
the microchip buffed and cleaned with a cotton swab applicator. The microchips were  
20 10 incubated with milli-Q water for about 30 minutes at room temperature before use.

The microchips were then washed with 50mM histidine buffer and biotinylated  
oligonucleotides (e.g., oligo dT12-btr) having a fluorophore in 50 mM histidine buffer  
25 were addressed to the capture sites using a standard A/C protocol (800 nAmps for 25  
seconds) to check the quality of the streptavidin microchips. The btr fluorophore was  
15 imaged using the appropriate excitation and emission filters for btr. The SDA primers  
(Seq. I.D. Nos. 20 and 21) were addressed to selected capture sites using the same standard  
30 A/C protocol.

As shown in Fig. 13, SDA is carried out at capture sites. Following denaturation of  
the double stranded target species, single stranded target molecules (e.g., a plus strand 10+  
20 shown in Fig. 13) are first addressed to the capture sites. For electronic hybridization of  
the various templates, double stranded DNA target sequence was first denatured at 95°C  
35 and mixed with an equal volume of 100 mM histidine buffer. The template mixture was  
then electronically hybridized to the capture SDA primers using a standard A/C protocol  
for hybridization (1.6  $\mu$ Amps for 60 seconds). After hybridization of the template  
40 25 mixture, the microchips were washed twice with water and incubated with 1 mg/ml BSA  
for 30 minutes at room temperature to block any non-specific binding sites. The  
microchips were washed again with water twice and pre-warmed at 60°C for 5 minutes.  
45 All SDA solutions were also pre-warmed at 60°C for 5 minutes. After pre-warming, the  
water was removed and the microchips were incubated with 10 $\mu$ l SDA reaction mixture  
30 (40mM K<sub>2</sub>HPO<sub>4</sub> pH 7.6, 1.6 mM each dCTP $\alpha$ S, dTTP, dATP, and dGTP, 8.3 mM  
50 MgCl<sub>2</sub>, 1.3 units BsoBI and 0.5 units Bst polymerase) for 30 minutes at 60°C in a

humidifying chamber. The reaction was stopped by removing the supernatant from the microchip surface to an eppendorf tube with 2 $\mu$ l of 100mM EDTA.

As indicated in Fig. 13, strand extension of the target nucleic acid of both plus and minus strands undergo strand displacement to form plus and minus single stranded amplicons (e.g., 12- and 13+). The plus and minus strand amplicons may each be electronically hybridized to adjacent or nearby unused primer pair sets.

In the instant example, following the SDA reaction, the microchips were washed three times with 0.5X SSC, pH 7.2. The SDA products were then denatured on the microchip *in situ* with addition of 0.5X SSC, pH 12.0 for 4 minutes in which the microchips were washed with fresh buffer every minute. The microchips were then washed with 0.5X SSC, pH 7.2 at least 3 times, with 4X SSC, pH 7.2 about 3 times. The microchips were then incubated with a 1  $\mu$ M mixture of btr-labeled reporter oligonucleotides in 4X SSC for 3 minutes at room temperature followed by extensive washing with 4X SSC at room temperature, then imaged with the appropriate laser and excitation/emission filters.

Although for simplicity in showing the efficiency of anchored SDA, this example carries out detection of SDA products following amplification, detection may be carried out during amplification using labeled target specific probes that are blocked at their respective 3' ends such as by incorporating a 3' phosphate group rather than a 3' OH on the terminus of the probe. Such labeled probes may further comprise single stranded nucleic acids which may be electronically addressed to the capture sites allowing detection of increasing signal as target and amplicon species are amplified at the capture pad site without the probe itself taking part in the SDA extension or amplification process.

In addition to the electronically controlled anchored SDA described above, two additional protocols were followed as controls wherein target nucleic acids were captured by passive hybridization followed by anchored SDA, and where SDA was carried out in solution. First, in the *passive* hybridization experiments, double stranded target nucleic acids were first denatured at 95°C for 5 minutes. The solution was then brought to a 4X SSC concentration with a 20X SSC (3M NaCl, 0.3 M NaCitrate) stock and 20ul of the mixture was pipetted onto the microchip (which had been previously electronically addressed with SDA primers) and incubated at room temperature overnight. Following the



target hybridization to the primers, SDA experiments were carried out as described above.

Second, where SDA was carried out in solution, no microchips were used. The reason for this is that the purpose of conducting solution based SDA was to compare the capacity to amplify target species in a multiplex format in solution versus on a microchip. The solution based SDA experiments were carried out in eppendorf tubes in a total of 50 $\mu$ l of SDA mix as described above.

In a first method of this example three different target nucleic acid species were amplified by SDA using primer pairs that were addressed to specific locations on an electronically addressable microchip. Ultra pure human placental DNA, *Chlamydia* genomic template and deoxynucleoside triphosphates were obtained from Becton Dickinson. Target templates for nucleic acids directed to detect the presence of gene sequence associated with hemochromatosis and Factor V were obtained using SDA bumper primers (Seq. I.D. Nos 22 and 23) and human placental DNA. PCR reaction conditions for amplifying such templates is well known to one of ordinary skill in the art of amplification. SDA capture primer pairs, bumpers, and signal probes for each test target species were synthesized and PAGE-purified by Oligos, Etc. (Oregon). The restriction site encoded into the primer sequences was BsoBI.

The following is a list of the various SDA primers and signal probes for each of the target species:

SDA primer biofacV10sSDA.213, (SEQ ID NO. 20)

5' [biot] ACCGCATCGAATGCATGTCCTCGGGTCTCTGGGCTAATAGGA 3'

SDA primer biofacVaSDA.297, (SEQ ID NO. 21)

5' [biot] ACGATT CAGCTCCAGACTTCTCGGGTCAGAATTTCTGAAAGG 3'

bumper primer facV10s.179, (SEQ ID NO. 22) 5' ACTACAGTGACGTGGACATC 3'

bumper primer facV10a.-127 (SEQ ID NO. 23) 5' TGTTATCACACTGGTGCTAA 3'

Signal probe facV10a.276 (SEQ ID NO. 24) 5' [BTR] CTGTATTCCTCGCCTGTC 3'

SDA primer chlaAL1.4811, (SEQ ID NO. 25)

5' [biot] CACGTAGTCAATGCATGTCCTCGGGTACAACATCAACACCTG 3'

SDA primer chlaAR1.4858, (SEQ ID NO. 26)

5' [biot] ACGATT CAGCTCCAGACTTCTCGGGTGAGACTGTAAAGATA 3'

bumper primer chlaBL1, (SEQ ID NO. 27) 5' CAGCAAATAATCCTTGG 3'

bumper primer chlaBR1, (SEQ ID NO. 28) 5' CATTGGTTGATGGATTATT 3'  
Signal probe chlaDIL.4826, (SEQ ID NO. 29) 5' [BTR] GTCGAGCCAAAATG 3'  
Signal probe chlaCP2.4841, (SEQ ID NO. 30) 5' [BTR]TTCCATCAGAAGCTGT 3'  
SDA primer haemsdas.6679, (SEQ ID NO. 31)  
5' [biot] CACGTAGTCAATGCATGTCCTCGGGTATAACCTTGGCTGTAC 3'  
SDA primer haemsdaa.6773, (SEQ ID NO. 32)  
5' [biot] ACGATTTCAGCTCCAGACTTCTCGGGTGCTCTCATCAGTCACA 3'  
bumper primer haempers.6596, (SEQ ID NO. 33) 5' TGAAGGATAAGCAGCCAAT 3'  
bumper primer haempcra.6773, (SEQ ID NO. 34) 5' CTCCTCTCAACCCCAATA 3'  
Signal probe haemreps.6712, (SEQ ID NO. 35) 5' [BTR]AGATATACGTGCCAGGTG 3'  
Signal probe haemreps.6733, (SEQ ID NO. 36) 5' [BTR]CTGATCCAGGCCTGGGTG 3'

As depicted in Fig. 16, biotinylated SDA primers for Factor V (FAC V),  
*Chlamydia* (CHL) and Hemochromatosis (HC) were anchored onto streptavidin-containing  
microchips and a mixture of Factor V, *Chlamydia* and Hemochromatosis templates were  
hybridized onto the primers electronically. Control template T12 was also anchored.

Anchored SDA was performed on microchips *in situ* at 60°C for 30 minutes as  
described previously and processed accordingly. As can be seen, no SDA amplicons can  
be detected when template is not hybridized to the SDA primers on the microchip (Fig.  
17). However, when a mixture of the templates are hybridized to the SDA primers,  
simultaneous amplification of the three amplicon systems can be seen (Figs. 18-20).  
Accordingly, when only one species of template is hybridized in the presence of all three  
types of SDA primer, only the area where the corresponding SDA primer is anchored  
shows a signal indicating amplification has taken place. This confirms the specificity, as  
well as the flexibility, of the anchored SDA system when done *in situ* on microchips.  
Interestingly, as shown in Fig. 21, when solution-based SDA is performed using the same  
three SDA primer sets, multiplex amplification is greatly compromised. Solution SDA  
was performed on Factor V, *Chlamydia* and Hemochromatosis separately, as well as  
together in one reaction (ALL) followed by analysis on a 6.0% non-denaturing  
polyacrylamide gel. As can be seen, all three systems amplify when done separately.  
However, when all three primer sets and templates are combined into one reaction, Factor  
V amplification is greatly depressed. Additionally, when the templates were hybridized to

5 the primers by passive hybridization, the amplification efficiency was significantly  
reduced, possibly due to the inefficient hybridization caused by template reannealing.  
10 These results underscore the need in the art for a system such as that of the current  
invention for a multiplex amplification system that can perform multiplex amplification  
5 and detection of target species without hindrance as may be observed in solution based  
and/or passive hybridization systems.

15 In a second embodiment of this example, the preparation of branched SDA target  
capture primer pairs may be synthesized by numerous means. In a preferred embodiment,  
the branched moiety may be produced as described below. First, as depicted in Fig. 22, the  
20 starting substrate for Y-primer synthesis is a biotin-conjugated lysine with a tert-butyloxy  
carbonyl-protected  $\alpha$ -amino terminal. The tert-butyloxy carbonyl (TBC) moiety on the  $\alpha$ -  
amino terminal allows selective attachment of the SDA primer arms separately. In this  
case, the  $\alpha$ -amino terminal is protected but the  $\alpha$ -amino terminal can react with carboxylic  
25 acid, allowing the SDA sense primer to be attached to the  $\alpha$ -amino terminal end. The  $\alpha$ -  
amino terminal end can then be deprotected with tri-fluoroacetic acid/dichloromethane  
(TFA/DCM), which removes the tert-butyloxy carbonyl moiety and allows attachment of  
the SDA antisense primer via the carboxylic acid terminal. This attachment sequence  
30 allows the formation of a Y-primer where both SDA primers are addressed to the branched  
moiety at their respective 5' ends. The resulting Y-shaped primer pair can then be  
20 attached to the streptavidin permeation layer on the microchip.

35 The synthesis of Y-shaped primer pairs for anchored SDA is intended to increase  
the overall efficiency of the SDA reaction twofold: 1) by placing the SDA primers in  
relatively close proximity of each other, thereby increasing the rate of interaction between  
40 extended amplicons of one strand and subsequent binding of the cleaved amplicon to the  
opposite strand primer; and 2) by increasing the density of primers in a given area over  
25 conventional oligonucleotide SDA primers. In the synthesis protocol above, the Y-primer  
is attached to the microchip permeation layer via a streptavidin-biotin bond; however,  
45 other amide-bond attachment chemistries can be used, including but not limited to prolinx,  
R-SH, or any other functional group onto the macromolecule. The branched primer pairs  
30 may be used in carrying out SDA reactions as described above.

#### 50 EXAMPLE 8

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

Still another example provides an asymmetric amplification method to address the problem of hybridization between sense and antisense amplicons that are generated during SDA. When using SDA, generally, both sense and anti-sense strands are generated in equal amounts. Under typical conditions of amplification, the complementary strands hybridize together. However, hybridization of oligonucleotides to specific sites on a microelectronic array (both for hybridization of amplimers to capture oligonucleotides and detection of hybridized material by fluorescently labeled reporter oligonucleotides) requires generation of single stranded species from the amplicons. Therefore, the complementary strands that are hybridized together must be separated prior to hybridization to capture upon the array and/or prior to detection by labeled reporter unless one strand is amplified more than the other (*i.e.* unless amplification is asymmetric). This is conventionally done using heat or chemical denaturation before or after electronic addressing. Asymmetric amplification removes the need for such thermocycling step.

15  
20  
25  
30  
35  
40  
45  
50  
55

A key feature of asymmetric amplification is the generation of a preponderance of one amplicon over its complementary amplicon sequence. In a solution environment, this method is typically accomplished by having a disproportionate ratio of amplifying primers. In the initial stages of the amplification process, the effective concentration of the sense and antisense amplifying primers being in large excess to template produces an environment conducive to exponential amplification of the original double stranded template material. As the reaction proceeds, the amplifying primer originally present in lesser amounts is effectively exhausted thereby leading to conditions of linear amplification by the primer remaining in excess. The particular effect of the polymerase mediated displacement of amplified material during SDA ensures that this linearly amplified material is free in solution and available for hybridization without the necessity for denaturation of double stranded species. With respect to objects of the invention, an alternative approach is to place both primers in solution at the same concentration, but to add a competitor that partially inhibits, or "poisons" generation of one strand. Over time this will also lead to a preponderance of one strand of the amplified target.

30  
35  
40  
45  
50  
55

Where capture probes are anchored, creation of predominantly one strand can be enhanced by designing anchored capture probes that are complementary to one strand of the amplicons being generated and released free into solution. In a preferred embodiment,

5 the capture probes are different from normal SDA primers in two respects. First, they  
preferably do not possess a functional restriction site, thereby blocking the endonuclease  
10 nicking/polymerase extension-displacement steps. Second, the 3' ends of the capture  
probes preferably are not suitable for extension by polymerase activity. During SDA this  
5 modified capture primers will hybridize to amplicon strands effectively pulling them out of  
the SDA pathway so that they will not be available for further amplification. The capture  
15 of such single strands may be directed to occur at a capture site located at a remote  
position from the site where SDA is occurring. Thus, a bias in the strandedness of the  
amplicon population will be generated, which is an effective form of asymmetric  
20 amplification due to limiting the quantities of one strand of amplification product.

In another example asymmetric amplification may be enhanced by including in the  
SDA reaction a competitive inhibitor of one of the primers of a given set of primers. As in  
the above example, the competitive primer is preferably either non-extendible or non-  
25 cleavable. The inclusion of the competitive primer biases the reaction toward the creation  
15 of single-strands through a linear reaction process.

Oligonucleotide sequences are rendered non-extendible using various means  
30 including blocking the 3' OH end, and mismatching the 3' terminal nucleotides(s) with  
respect to the template sequence. Oligonucleotide sequences are rendered non-cleavable  
by modifying the oligonucleotide backbone through the inclusion of modified linkages  
20 such as phosphorothioates or more simply by changing the sequence at the restriction  
endonuclease recognition site. Probes modified as such remain fully competent for  
35 hybridization. The sequence of the competitor is preferably identical to (or nearly identical  
to) that of one of the amplification primers. The competitor can therefore compete with  
the amplification primer for hybridization with a target sequence. When bound to the  
40 target sequence, the competitor either (1) cannot be extended by DNA polymerase, or (2)  
25 can be extended to produce a copy of the target sequence. In the case where the  
competitor is extended, the competitor is modified such that resultant copies of the target  
sequence cannot be cleaved by a restriction enzyme. Different types of competitors are  
45 used depending on the amplification method being used.

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In PCR, the competitor is modified such that it cannot be extended. Appropriate modifications are described above. In each cycle of the reaction, the competitor will compete with one of the PCR primers for hybridization to available target sequences. For example, in a reaction where the competitor is added at 10% the concentration of the PCR primer, roughly 10% of hybridization events will be abortive in that an extension product cannot be produced. The opposite PCR primer is free to hybridize to all available target sequences and be extended. Therefore, a bias of about 10% in the relative number of the two extension products is produced in any given cycle. While a 10% bias in early cycles may not be significant since target concentration is low, such a bias will produce a high concentration of single-stranded material in late stage cycles (where nM quantities or greater of the extension products are being produced).

10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In SDA, several methods are preferable. Use of a non-extendible competitor will bias the production of double-stranded templates which will allow the nicking and extension reaction to preferentially produce one of the single-stranded displaced products. Use of an extendible, non-cleavable competitor leads to asymmetry by creating double-stranded products that cannot participate in the nicking/displacement reaction. Use of both types of competitors may be optimal as extension products produced from the non-cleavable primer become part of double-stranded molecules when only one strand can be nicked and displaced. (see Fig. 14).

#### EXAMPLE 9

In this example of the invention, SDA is carried out in conjunction with an electronically addressable microchip wherein the atmospheric pressure of the SDA reaction is elevated.

Where genomic nucleic acid is used, it is preferred that it be cleaved into fragments of between approximately 250-500 bp. This may be done by a restriction enzyme such as HhaI, FokI or DpnI. The selection of the enzyme and the length of the sequence should be such that the target sequence sought will be contained in its entirety within the generated fragments or that at least a sufficient portion of the target sequence will be present in the fragment to provide sufficient binding of SDA amplification primers. Other methods for generating fragments include PCR and sonication.

5 The primers used in this method generally have a length of 25-100 nucleotides.  
Primers of approximately 40 nucleotides are preferred. The primer nucleic acid sequence  
10 should be substantially homologous to the target sequence such that under high stringency conditions hybridization between primer and template nucleic acid will occur.

5 Target nucleic acid fragments are denatured to render them single stranded so as to permit binding of the primers to the target strands. Raising the temperature of the reaction  
15 to approximately 95°C is a preferred method for denaturing the nucleic acids. Other methods include raising pH; however, this will require lowering the pH in order to allow the primers to bind to the target. Following the formation of single stranded target  
20 10 molecules, SDA is performed as discussed in the numerous examples discussed herein. Typically, the SDA reaction includes the use of at least one substituted nucleotide during primer extension to facilitate nicking of one strand during amplification. The nuclease may be any nuclease typically useful for SDA as discussed earlier.

25 In a preferred embodiment of this method, atmospheric pressure is elevated either  
15 before or after all the SDA reaction components are combined. The pressure is elevated to reduce star activity to effectively enhance the specificity of the restriction endonuclease for its target. The application of elevated pressure may also increase the specificity of primer  
30 interaction with the template nucleic acid and the overall rate of reaction of the enzymes employed, thereby reducing the time required for the SDA reaction while increasing its  
20 specificity. By reducing star activity, template independent amplification is decreased thereby reducing the competitive consumption of reagents by non-specific amplification.

35 Elevated pressure can be supplied during the amplification by various methods. For example, the reactions could be run in high pressure vessels. The reactions may also  
40 be run by placing the container in a reaction chamber attached to or part of a high-pressure apparatus (High Pressure Equipment Co., Erie, PA). It may be advantageous to overlay  
25 the aqueous reaction media with an immiscible phase, such as silicon oil (Sigma) by which pressure can be applied to the aqueous solution containing the target nucleic acid,  
45 nucleosidetriphosphates, and enzymes. Preferably, the pressure is elevated in the range of about 100 to about 500 atmospheres.

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

Polymerases useful in this method include those that will initiate 5'-3' polymerization at a nick site. The polymerase should also displace the polymerized strand downstream from the nick, and, importantly, should also lack any 5'→3' exonuclease activity and be heat stable. Polymerases, such as the large fragment of DNA polymerase I and the exonuclease deficient Klenow fragment of DNA polymerase I and a similar fragment from the Bst polymerase (New England Biochemicals, Beverly, Mass.) are useful. SEQUENASE 1.0 and SEQUENASE 2.0 (U.S. Biochemical), T5 DNA polymerase, and Phi29 DNA polymerases are also useful. Generally, thermophilic DNA polymerases are preferred. The exonuclease deficient thermophilic Klenow fragment of Bst DNA polymerase from *Bacillus stearothermophilus* (New England Biochemicals, Beverly, Mass.) is most preferred.

In this method, a single reaction temperature may be employed after denaturation has occurred, and such temperature should be high enough to set a level of stringency that minimizes non-specific binding but low enough to allow specific hybridization to the target strand. In addition, use of temperature preferably from about 45°C. to about 60°C should support efficient enzyme activity. Denaturation of the enzymes and nucleic acid is to be avoided.

During the SDA reaction cycles, theoretically about 20 repetitions or cycles will result in about a  $10^6$ -fold amplification (*i.e.*,  $SDA\ X2^{20}=10^6$ ). Typically,  $10^8$ -fold or greater amplification is seen in about 30 minutes of amplification.

High pressure SDA is beneficial for various uses including generation of high fidelity single-stranded nucleic acid probes or single-stranded templates for sequencing. Toward this goal, high pressure SDA can be conducted either with a single primer or using two primers wherein one primer is in excess over the other. The result is excess production of one displaced single strand over the other.

The presence of the amplified target then can be detected by any number of methods. One method is to detect reaction products of a specific size by means of gel electrophoresis. This method is particularly useful when the nucleotides used are labeled with a radio-label, such as  $^{32}P$ . Other methods include labeling the nucleotides with a physical label, such as biotin. Biotin-containing reaction products can then be identified by means of avidin bound to a signal generating enzyme, such as peroxidase. Another



method is elongation of a fluorescently labeled internal primer.

Detection systems useful in the practice of this invention comprise homogeneous systems, which do not require separation, and heterogeneous systems. In each system, one or more detectable markers are used and the reaction or emission from the detection system is monitored, preferably by automated means. Examples of homogeneous systems include fluorescence polarization, enzyme mediated immunoassays, fluorescence energy transfer, hybridization protection (e.g., acridinium luminescence) and cloned enzyme donor immunoassays. Examples of heterogeneous systems include enzyme labels (such as peroxidase, alkaline phosphatase and beta-galactosidase), fluorescent labels (such as enzymatic labels and direct fluorescence labels (e.g., fluorescein and rhodamine)), chemiluminescence and bioluminescence. Liposomes or other sac like particles also can be filled with dyes and other detectable markers and used in such detection systems. In these systems, the detectable markers can be conjugated directly or indirectly to a capture moiety or the amplified products can be generated in the presence of a receptor which can be recognized by a ligand for the receptor.

#### Protocol for Strand-Displacement Amplification (SDA) Under Elevated Pressure

Amplification reactions utilize approximately 100ng of genomic DNA (Factor V) in a total volume of 50  $\mu$ l. The genomic DNA (human placental DNA; Becton-Dickinson) is denatured at 95°C for 5 minutes followed by centrifugation to collect condensate. Next, 1  $\mu$ l of SDA primer mix is added (50  $\mu$ M each reaction) and incubated at 60°C for 5 minutes. SDA mix (40 mM  $k_2HPO_4$  pH 7.6, 1.4 mM each dCTP $\alpha$ S, dTTP, dATP and dGTP, 8.3 mM  $MgCl_2$ , 40 units/rxn BsoBI (New England Biochemicals), 15.6 units/rxn Bst polymerase (New England Biochemicals), and 0.05  $\mu$ M each SDA bumper primers are added and pre-warmed for 5 minutes at 60°C followed by addition of the mix to SDA primers and target sample. Silicon oil is added to the top of the reaction tubes and placed in high a pressure chamber. The pressure is elevated to between 100 and 500 atmospheres and incubate at 60°C for 30 minutes. Following the reaction period, the pressure is reduced to atmospheric pressure and stopped by addition of 10  $\mu$ l of 100 mM EDTA. SDA products are visualized by electrophoresing on 6% non-denaturing polyacrylamide gels. The gels are stained with ethidium bromide and photographed under UV-

5  
fluorescence.

10 Alternatively, it is possible to use a device wherein the temperature and/or pressure is elevated prior to the addition of the polymerases and/or restriction endonuclease.

5 The use of elevated pressure can also be used in the performance of anchored SDA, or any SDA procedure as described above. Specifically, when anchored SDA is performed on electronically addressable microchips, elevated pressure should decrease star activity and increase efficiency by reducing primer independent amplification.

#### EXAMPLE 10

20 10 In another example, SDA may be used in conjunction with electronically addressable microchips wherein the SDA reaction is "ligation-dependent" or "ligation-based". This method involves the SDA amplification of a ligated probe using a pair of universal amplification primers. The amplification primers are universal in the sense that they are designed to amplify all ligated probes in a test reaction whether the reaction is  
25 15 multiplexed or directed to a singular target. The ligated probe is formed by ligating together a pair of ligation probes that have hybridized to a target sequence. No bumper primers are necessary.

30 In another embodiment, a method of ligation-based SDA is provided where the method is unassisted by an electronic microchip. In this embodiment it is not necessary to, *inter alia*, anchor any primers, which is a procedure that assists in separating primer sets  
35 20 during multiplex amplification, because the primers are universal -- there is no need to direct target sequences to specific primers.

40 The following functional descriptions of the oligonucleotide reagents are not intended to define or limit their actual physical composition. Rather, the description  
25 merely demonstrates that each reagent exhibits certain functional characteristics. Thus, it should be noted that the functional regions of a given oligonucleotide reagent may overlap, or in fact be co-extensive, as where a specific nucleic acid sequence is able to accomplish  
45 more than one function. Additionally, the individual base sequence in any given oligomer depends upon the target nucleic acid of interest, the restriction enzyme chosen for use in  
30 SDA, or an arbitrary sequence chosen for portions of the amplification primers and ligation probes so that a degree of universality can be incorporated into the amplification  
50

5 protocol.

10 In operation, as illustrated in Figure 23(a-c), the ligation-based SDA method uses a pair of ligation probes that anneal to adjacent nucleic acid sequences on a target.

5 Functionally, the pair of ligation probes bind to a target nucleic acid sequence such that they can be ligated together while they are annealed to the target to form a ligated probe template. Ligation will occur only following hybridization of both ligation probes of a  
15 ligation probe pair to a target sequence.

20 The first *ligation* probe can be divided into three functional regions: a 5' region able to hybridize to target nucleic acid; a middle region; and a 3' region comprising a nucleic acid sequence that is able to hybridize to the first *amplification* primer. The second  
25 *ligation* probe can also be divided into three functional regions: a 5' region having a nucleic acid sequence identical to nucleic acid sequences found in the second *amplification* primer and having a restriction endonuclease recognition site; a middle  
30 region; and, a 3' region able to hybridize to target nucleic acid.

15 With respect to the amplification primers, the first amplification primer can be divided into two functional regions: a 5' region containing a restriction endonuclease recognition site and a 3' region that is able to hybridize to the first ligation probe. The  
30 second amplification primer can also be divided into two functional regions: a 5' region that contains a recognition site for a DNA restriction endonuclease and a 3' region  
35 comprising nucleic acid sequence having the same sequence as the 5' region of the second ligation probe.

40 The ligation-based SDA reaction comprises a number of component steps. In Step 1, the pair of ligation probes anneal to adjacent sequences of single-stranded target nucleic acid such that the second ligation probe hybridizes to the target strand at a position on the  
45 target that is 3' to the hybridization position of the first ligation probe. In Step 2, DNA ligase catalyzes the ligation of the two ligation probes to form the ligated probe template. In a preferred embodiment, the 3' end of the ligated probe template is modified to prevent  
50 primer extension from that end (Figure 23(a-c)).

55 In Step 3, the first amplification primer binds to the 3' end of the ligated probe template such that the amplification primers extend beyond the end of the template forming a 5' overhang. In a preferred embodiment, DNA polymerase catalyzes new DNA

5 synthesis from the 3' end of the first amplification primer causing the ligated probe to be  
displaced from the target nucleic acid. This results in the release of single-stranded target  
10 nucleic acid and the creation of double-stranded DNA oligonucleotide having a 5'  
overhang (labeled Product I, Fig. 23). The release of single-stranded target nucleic acid  
5 and the creation of the double-stranded oligonucleotide occurs without the assistance of  
bumper primers. Moreover, the target single strand becomes available for further binding  
15 of unligated first and second ligation probes.

Product I thus comprises a first strand having a sequence from 5' to 3'  
20 corresponding to the ligated probe template, and a second strand complementary to the  
ligated probe template strand with an additional nucleic acid sequence at its 5' end  
corresponding to the 5' end of the first amplification primer. This double stranded DNA  
molecule is capable of undergoing a series of SDA reactions that produce single stranded  
25 DNA molecules able to be bound and amplified by the universal amplification primers.  
The double stranded DNA molecules that result from these reactions are also susceptible  
15 to amplification. Nicking by a restriction endonuclease, followed by primer extension and  
strand displacement, substantially regenerates the double stranded DNA starting material.  
Together, these ligation-dependent SDA reactions ultimately amplify oligonucleotide  
30 sequences corresponding to the ligated probe, thereby allowing the detection of the target  
sequence. These reactions are described in detail below.

20 In Step 5, Product I is nicked by a restriction enzyme to create Product II. In Step 6,  
35 Product II undergoes primer extension and strand displacement from the nick, resulting in  
Product III and Product IV. Product III is essentially the same as Product I except that the  
first strand of Product III (which corresponds to the first strand of Product I) contains an  
40 additional sequence at its 3' end complementary to the 5' end of the first amplification  
25 primer. Product IV is a single-stranded molecule with a sequence comprising the first  
strand of Product II located 3' to where this strand was nicked by the restriction  
endonuclease.

45 In Step 7, Product III is nicked by a restriction endonuclease to create Product V.  
In Step 8, Product V undergoes primer extension and strand displacement to create Product  
30 VI and Product VII. Product VI is essentially the same as Product III. Product VII is a  
single stranded DNA molecule comprising the nicked strand of Product V located 3' to the  
50

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

nick site.

In Step 9, the second amplification primer binds to Product VII. In Step 10, Product VII undergoes a primer extension reaction in both directions to create Product VIII. Product VIII is a double stranded nucleic acid molecule, the first strand having a sequence corresponding to product VII plus an additional 3' sequence that is complementary to the 5' region of the second amplification primer, and a second strand that is complementary to the first strand. In Step 11, Product VIII is nicked with a restriction endonuclease to create Product IX. Product IX is essentially the same as Product VIII except that the 5' end of Product IX contains a nick in the nucleic acid corresponding to the 5' region of the second amplification primer. In Step 12, Product IX undergoes primer extension and strand displacement to create Products X and XI. Product X is the same as Product VIII. Product XI is a single stranded nucleic acid molecule with a sequence corresponding to the sequence 3' of the nick, on the nicked strand of Product IX. In Step 13, Product XI is bound by the first amplification primer and in step 14, primer extension in both directions results in Product XII. Product XII is a double stranded nucleic acid molecule similar to Product III in the sense that it can enter the above described reaction pathway following step 6 and prior to step 7. Thus, an initial reaction product of the ligation-dependent SDA pathway is ultimately substantially regenerated.

As described earlier, the SDA reaction may be carried out using anchored probes. With regard to ligation-based SDA, the anchored probes are preferably either one or both of the amplification primers or one or both of the ligation probes.

#### Experimental data for EXAMPLE 10

##### Experiment 1

In this example, a general protocol for the preferred ligation-based SDA of a target nucleic acid is provided. Concentrations and volumes of reaction components, and time and temperature profiles may be adjusted as necessary. Volumes assume a 25 $\mu$ l ligation reaction volume and a 50 $\mu$ l final reaction volume for SDA.

In a 250 $\mu$ l microcentrifuge tube, a 5 $\mu$ l aliquot of an aqueous ligation probe solution is added such that the final concentration of each probe in a 25 $\mu$ l ligation reaction volume will be 5nM. Next, 10 $\mu$ l of a solution of non-specific (carrier) DNA (*e.g.*, Calf thymus

DNA is added to a final concentration of 20-100 $\mu$ l/ml. Next, 5 $\mu$ l of the sample containing the template nucleic acid (e.g. Cell lysate or purified genomic DNA) at an appropriate concentration is added and the tube is placed at 60°C for 3 minutes to allow temperature equilibration. Following equilibration, 5 $\mu$ l of a solution containing a thermostable DNA ligase is added along with sufficient 5X strength mixture of buffer components necessary to allow function of the DNA ligase, and to allow probe hybridization. See Table IV.

The 25 $\mu$ l ligation reaction is incubated at 60°C for 15 minutes and then 20 $\mu$ l of an SDA stock mix containing additional buffer components, dNTPs, and amplification primers, is added to give final reaction concentrations (in 50 $\mu$ l) as shown in Table V. In one embodiment an additional step is included where the reaction is heated to 95°C for 3 minutes to denature the ligated probes from the template and then the tube is equilibrated at 60°C for 3 minutes. To this reaction mixture 5 $\mu$ l of liquid containing the SDA enzymes is added to give the following final concentrations in a 50 $\mu$ l final reaction volume:

*BSOB1* restriction enzyme: 0.8 enzyme units/ $\mu$ l (40U/rxn)

*Bst* DNA polymerase: 0.32 enzyme units/ $\mu$ l (16 U/rxn)

The reaction mixture is incubated at 60°C for 30 minutes then the reaction is stopped by placing the reaction mixture on ice.

Table IV

Buffer Component	Final Concentration in Reaction for Each Ligase	
	<i>Taq</i> DNA ligase (1 U/rxn)	<i>Pfu</i> DNA ligase (0.2 U/rxn)
Tris-HCl pH 7.6	10 mM	10 mM
Potassium Acetate	25 mM	25 mM
Magnesium Acetate	10 mM	10 mM
Dithiothreitol	1 mM	1 mM
Nicotinamide adenine dinucleotide	1 mM	NONE
Adenosine triphosphate	NONE	10 $\mu$ M

Table V

SDA Component	Final Concentrations in 50 $\mu$ L Reaction (Note: includes contribution from ligation reaction)
Potassium phosphate	35 mM
Bovine serum albumin	80 $\mu$ g/ml
Magnesium acetate	10 mM
Deoxynucleotide triphosphates (equal mixture of dATP, dC <sub>as</sub> TP, dGTP, TTP)	1.4 mM
Amplification primers (S1 and S2)	250 nM

## Experiment 2

In this further example, the *Salmonella* spaQ gene (a portion of which is indicated on Fig. 23 d and designated SEQ. ID. No. 41) potentially present in a sample is amplified. The reaction protocol as described in Experiment 1 is followed using the ligation probes LP1 (SEQ. ID. No. 37) and LP2 (SEQ. ID. No. 38) and amplification primers S1 (SEQ. ID. No. 39) and S2 (SEQ. ID. No. 40) which are illustrated in Figure 23(d). The example described in Experiment 2 is intended to have general applicability. One could create different target-specific ligation probes for use with the amplification primers S1 and S2 by replacing the sequences of ligation probes L1 and L2 complementary to the spaQ gene with sequences complementary to another target nucleic acid of interest. Moreover, amplification primers S1 and S2, such as those depicted in Fig. 23(d) may be used in a multiplex amplification of more than one target nucleic acid.

## Experiment 3

At high concentrations of ligation probe, ligase may catalyze the ligation of the ligation probes in a target-independent manner. The resulting ligated probe can support SDA and may thus create a false positive signal. In this further example, a preferred aspect of ligation-dependent SDA is described where this problem is overcome by rendering the ligation probes initially incapable of being ligated together by ligase. In this embodiment, a

5 pair of unligatable probes is rendered ligatable to allow target-specific, ligation-dependent SDA.

10 Generally, the amplification of a background molecule that is target independent may be prevented by modifying the ends of the ligation probes that are involved within the  
5 ligation junction. This can take place in several ways. One such modification involves the modification (including removal, blocking, *etc.*) of the 3' hydroxyl group present on the 3'  
15 terminal nucleotide of the second ligation probe (the upstream probe). Another such modification involves the modification (including removal, blocking, *etc.*) of the 5' phosphate group present on the 5' terminal nucleotide of the first ligation probe (the  
20 downstream probe). Various methods have been and can be devised wherein the removal and or alteration of these modifications occurs preferentially in the presence of target DNA.

25 Specifically, one aspect of this example provides for modifying the 3' hydroxyl group present on the 3' terminal nucleotide of the second ligation probe (the upstream  
15 probe) to prevent blunt end ligation between the ligation probes. The modified unligatable probe is rendered ligation competent using an endonuclease, preferably Endonuclease IV. This reagent is able to excise 3' terminal nucleotides from  
30 oligonucleotides and thus is used to excise the 3' terminal nucleotide of the second ligation probe to reveal a new 3' terminal nucleotide with a 3' hydroxyl group. This reaction is  
20 more preferred when the ligation probe substrate is associated with target DNA and less preferred when the ligation probe substrate is unassociated with other DNA molecules.  
35 Consequently, once the ligation probes are bound to target DNA, the endonuclease (preferably Endonuclease IV) is able to excise the 3' terminal nucleotide of the second ligation probe to reveal a new 3' terminal nucleotide with a 3' hydroxyl group. The free 3'  
40 hydroxyl group of the second ligation probe, along with the free 5' phosphate group of the  
25 first ligation probe, are now substrates for ligation by DNA ligase.

45 Since endonuclease tends to operate more efficiently when the substrate oligonucleotide is double stranded it will preferentially excise the 3' terminal nucleotide of the second ligation probe when this probe is bound to target DNA, not when it is free in  
30 solution. Because the endonuclease preferentially renders the initially ligation-incompetent ligation probes ligation-competent when they are in the presence of target DNA, the target  
50



independent amplification of background molecules is decreased.

Another aspect of this example provides for the modification (including removal, blocking, *etc.*) of the 5' phosphate group present on the 5' terminal nucleotide of the first ligation probe (the downstream probe) to prevent blunt end ligation between the ligation probes. The modified unligatable probe is rendered ligation-competent using a DNA polymerase with exonuclease activity. This reagent will allow DNA polymerization (new DNA synthesis) to occur from the 3' end of the upstream probe (the second ligation probe) into the 5' end of the downstream (first ligation) probe. When the polymerase contacts the 5' end of the first ligation probe it will begin to excise nucleotides from the 5' end. As it excises nucleotides from the first ligation probe, nucleotides are added to the 3' end of the second ligation probe. In essence, this moves the "gap" between the first and second ligation probes, the junction to be ligated by ligase, from 5' to 3'. By controlling the amount and/or type of free nucleotide present in solution, the degree of excision and replacement can be limited. Following dissociation of the polymerase the junction contains a free 3' hydroxyl group and a free 5' phosphate group, both of which are substrates for ligation by DNA ligase. As indicated above, this reaction is more preferred when the ligation probe substrate is associated with target DNA and less preferred when the ligation probe substrate is unassociated with other DNA molecules. Again, this is because the reaction that renders the ligation probes ligatable prefers that the ligation probes be annealed forming dsDNA. As is understandable to one skilled in the art, such annealing is preferred for target DNA rather than annealing to non-target DNA. Thus, independent amplification of background molecules is decreased.

Yet another aspect of this example provides for blocking ligation using base-pairing mismatching. Here, ligation is prevented between the first and second ligation probes by having the 5' end of the downstream (first) probe contain one or more mismatched bases. If a probe is said to contain a mismatched base, it should be understood to mean that the probe contains a nucleotide that is not complementary to target DNA sequences, in a region of the probe otherwise complementary to the target DNA. Mismatched bases prevent ligation by DNA ligase until the mismatched bases are excised, as in the above stated example, with DNA polymerase.

To demonstrate the exonuclease/ligase-dependent SDA (XL-SDA) aspect of this invention, as described in this further example, the nine sets of ligation probes shown in Table VI were synthesized. These probes were designed to identify the various bacterial species shown. The probes have regions complementary to the specific bacterial genes and regions designed for SDA amplification primer binding.

Table VI

Bacterial gene, product	Genus/Species/serotype identified	Ligation probe 1 (5'-3')	Ligation probe 2 (5'-3')
<i>stx</i> <sub>1</sub> , Shiga-like toxin-I	Shiga toxin-producing <i>E. coli</i> (STEC) and <i>Shigella dysenteriae</i> type I	GAGGGCGGTTTAATAA TCTACGGTGGTCGAGT ACGCCTTAA (SEQ. ID. No. 45)	CGATTCCGCTCCAGACTT CTCGGGTGTACTGAGATC CCCTTGTCAGAGGGATAG ATCCAGAGG (SEQ. ID. No. 46)
<i>stx</i> <sub>2</sub> , Shiga-like toxin-II	STEC	GATGGAGTTCAGTGGT AATACAAATGTGGTCGA GTACGCCCTTAA (SEQ. ID. No. 47)	CGATTCCGCTCCAGACTT CTCGGGTGTACTGAGATC CCCTGGTTTCATCATATCT GGCGTT (SEQ. ID. No. 48)
<i>eaeA</i> , intimin	<i>E. coli</i> O157:H7	GACGCTGCTCACTAGA TGTCTAGGTCGAGTAC GCCTTAA (SEQ. ID. No. 49)	CGATTCCGCTCCAGACTT CTCGGGTGTACTGAGATC CCCTGGTTATAAGTGCTT GATACTCCAG (SEQ. ID. No. 50)
<i>spaQ</i> , surface antigen-presenting protein	<i>Salmonella</i> species	GATGATGTCATGTTGC AATGTCCTGGTCGAGT ACGCCTTAA (SEQ. ID. No. 51)	CGATTCCGCTCCAGACTT CTCGGGTGTACTGAGATC CCCTCATTTAACTATCCC GTCTCGT (SEQ. ID. No. 52)
<i>gnd</i> , 6-phosphogluconate	<i>Salmonella typhi</i> , <i>Salmonella</i>	GAGTAATTACCGTCTT CATCTTTTTTTGGTCGA	CGATTCCGCTCCAGACTT CTCGGGTGTACTGAGATC

dehydrogenase	<i>paratyphi</i>	GTACGCCTTAA (SEQ. ID. No. 53)	CCCTGGCTTCATCAAGAA TAACATCTATC (SEQ. ID. No. 54)
<i>ipaH</i> , pathogenicity- associated gene	Shigella species and enteroinvasive <i>E. coli</i>	GATTTACGGACTGGTT CTCCCTTGGTCGAGTA CGCCTTAA (SEQ. ID. No. 55)	CGATTCCGCTCCAGACTT CTCGGGTGTACTGAGATC CCCTTCAGAAGCCGTGAA GAGAATG (SEQ. ID. No. 56)
<i>sodB</i> , superoxide dismutase	Campylobacter species	GACCAAAACCATCCTG AACCATGGTCGAGTAC GCCTTAA (SEQ. ID. No. 57)	CGATTCCGCTCCAGACTT CTCGGGTGTACTGAGATC CCCTTTCTAGTTTTTGATT TTTAGTATTATA (SEQ. ID. No. 58)
<i>asd</i> , aspartate semialdehyde dehydrogenase	Vibrio species	GAGTAGAGGTATGTGA TGAGCCAATGGTCGAG TACGCCTTAA (SEQ. ID. No. 59)	CGATTCCGCTCCAGACTT CTCGGGTGTACTGAGATC CCCTCTTTGGCTAAACTC GGTTTTT (SEQ. ID. No. 60)
<i>lcrV</i> , Yersinia V- antigen	Yersinia species	GATTAGCTGAGCTTAC CGCCGTGGTCGAGTAC GCCTTAA (SEQ. ID. No. 61)	CGATTCCGCTCCAGACTT CTCGGGTGTACTGAGATC CCCTCCGTAGCAAGTTGC GTGAAG (SEQ. ID. No. 62)

The probes were added to identical sets of ligation-SDA reaction such that the number of ligation probe sets in the reactions increased in the order: 1 set (*spaQ*), 5 sets (*spaQ*, *stx1*, *stx2*, *sodB*, *ipaH*), 6 sets (as 5+*lcrV*), 7 sets (as 6+*asd*), 8 sets (as 7+*eaeA*), 9 sets (as 8+*gnd*), and such that the final concentration of each probe was 5nM.

A total extract of *Salmonella enteritidis* genomic DNA was added as a template such that the estimated number of genome equivalents was either  $10^5$ ,  $10^4$ ,  $10^3$  or zero as a negative control. XL-SDA reactions were performed as described below, and the reaction products analyzed by both acrylamide gel electrophoresis and electronic hybridization on a microelectrode array.

XL-SDA reactions were performed as follows although the concentrations and volumes of reaction components and time/temperature profiles may be adjusted as necessary. The volumes used assume a 25 $\mu$ l ligation reaction volume and a 50 $\mu$ l final reaction volume for SDA.

In a 250 $\mu$ l microcentrifuge tube, solutions of the following reagents were combined to give the final concentrations shown: (1) two (or more) target-specific ligation probes (e.g. probe exo-LP1 having a 5' sequence substantially complimentary to a portion of the target sequence of interest and a 3' sequence complimentary to a universal amplification primer and probe exo-LP2 having a 3' end sequence substantially complementary to a portion of the target sequence located downstream of LP1 and a 5' end sequence identical to a second universal amplification primer) to give probe concentrations of 5nM of each probe; and, (2) a solution containing the template DNA of interest.

The exonuclease/ligation reaction was initiated by the addition of the following: a thermostable DNA ligase (such as *Taq* DNA ligase or *Pfu* DNA ligase); a thermostable DNA polymerase having 5'-3' exonuclease activity, (such as *Taq* DNA polymerase); buffer salts to give final concentrations shown in Table VII below; and dATP at 2.8mM in a 25 $\mu$ l reaction.

Table VII

Buffer Component	Final concentration in reaction for each ligase	
	<i>Taq</i> DNA ligase (1 U/rxn)	<i>Pfu</i> DNA ligase (0.2 U/rxn)
Tris-HCl pH 7.6	10 mM	10 mM
Potassium Acetate	25 mM	25 mM
Magnesium Acetate	10 mM	10 mM
Dithiothreitol	1 mM	1 mM
Nicotinamide adenine dinucleotide	1 mM	NONE
Adenosine triphosphate	NONE	10 $\mu$ M

The ligation/exonuclease reaction was incubated at 60°C for 15-30 minutes. Then, 20µl of an SDA stock mix containing additional buffer components, a mixture of dNTPs such that the final reaction contains all four dNTPs, and amplification primers, is added to give the final reaction concentrations (in 50µl) shown in Table VIII.

Table VIII

SDA Component	Final Concentrations in 50µl reaction
Potassium phosphate	35 mM
Bovine serum albumin	80 µg/ml
Magnesium acetate	10 mM
Deoxynucleotide triphosphates (dGTP, dC <sub>as</sub> TP, TTP)	1.4 mM
Amplification primers (S1 and S2)	250 nM

Then, 5µl's of a solution containing the SDA enzymes is added to give the following final concentrations: *Bso*B1 restriction enzyme at 0.8 enzyme units/µl (40U/rxn) and *Bst* DNA polymerase at 0.32 enzyme units/µl (16 U/rxn). This reaction mixture is then incubated at 60°C for 30 minutes to allow the SDA reaction to proceed. The reaction is stopped by placing it on ice and the amplified products are detected.

The reaction products generated were analyzed by both acrylamide gel electrophoresis and electronic hybridization on a microelectrode array. An analysis of 5µl of the XL-SDA reactions by acrylamide gel electrophoresis demonstrated that specific amplification product is made in a template concentration-dependent manner in all combinations of ligation probes. To demonstrate specific amplification of the *Salmonella enteritidis spaQ* gene sequence, the ligation-SDA reaction products were analyzed on a microelectrode array where specific capture probes for five of the bacterial genes are pre-arranged at discrete locations. Figure 24 shows that in all samples analyzed, the *spaQ* sequence was detected.

5

10

15

20

25

30

35

40

45

50

55

The foregoing is intended to be illustrative of the embodiments of the present invention, and are not intended to limit the invention in any way. Numerous variations and modifications of the present invention may be effected without departing from the true spirit and scope of the invention. As is understandable to one of ordinary skill in the art, each of the embodiments as disclosed above may be used together in any combination. For example, SDA may be carried out in connection with an electronically addressable microchip wherein amplification primers specific for a target nucleic acid (such as branched or unbranched primer pairs having complementary sequence to ligation probes or other target nucleic acids of interest) are anchored to an electronically addressable capture pad, target nucleic acid is electronically addressed to such capture pads, and SDA is performed under high pressure. In another example, SDA may be carried out in connection with an electronically addressable microchip wherein allele-specific amplification primers (such as branched or unbranched primer pairs) are anchored to an electronically addressable capture pad, target nucleic acid is electronically addressed to such capture pads, and SDA is performed under high pressure or in the alternative at atmospheric pressure. In still another combination example, SDA may be carried out in connection with an electronically addressable microchip wherein the SDA reaction is carried out using noncleaveable primers or under asymmetric amplification conditions. Additionally, other combinations may include ligation-based SDA in combination with the electronically addressable microchip either under elevated or normal atmospheric pressures. As is understandable to one of ordinary skill in the art, many other combinations are possible.

Although the invention has been described with respect to specific modifications, the details thereof are not to be construed as limitations, for it will be apparent that various equivalents, changes and modifications may be resorted to without departing from the spirit and scope thereof and it is understood that such equivalent embodiments are to be included herein.

All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

## Claims

5

10

15

20

25

30

35

40

45

50

55

What is claimed is:

1. A method for the amplification, multiplex assaying, and detection of target nucleic acids of interest using a bioelectronic microchip comprising the steps of:
  - a. introducing at least one of said target nucleic acids of interest onto a bioelectronic microchip having a plurality of electronically addressable capture sites;
  - b. amplifying said target nucleic acids to form amplicons of said target nucleic acids wherein said target nucleic acids are amplified using an amplification reaction wherein a multiplicity of sets of amplification primers are used that include unequal ratios of upstream and downstream primers;
  - c. capturing said amplicons of said target nucleic acids onto specified capture sites of said microchip by a capture probe that has specificity for said amplicons; and
  - d. detecting the presence of said captured amplicons.
2. The method of claim 1 wherein amplification of the target nucleic acid is by strand displacement amplification.
3. The method of claim 1 wherein amplification of the target nucleic acid is by at least one of the methods selected from the group consisting of strand displacement amplification, allele-specific strand displacement amplification, nucleic acid sequence-based amplification, and ligation-based strand displacement amplification.
4. The method of claim 1 wherein said capturing of amplicons in (c) is carried out by passive hybridization.
5. The method of claim 1 wherein said capturing of amplicons in (c) is carried out by electronic biasing of said capture sites.



5

6. The method of claim 1 wherein said amplification is carried out in part using noncleavable and/or non-extending primers.

10

5

7. The method of claim 1 wherein said amplification, multiplex assaying and detecting are carried out either consecutively or simultaneously in relation to one another.

15

8. The method of claim 1 wherein detection of amplicons is by at least one of fluorescence, chemiluminescence, and electrochemiluminescence.

20

10

9. The method of claim 1 wherein target nucleic acids of interest are derived from a source selected from the group consisting of blood, sputum, stool, tissue, media, semen, hair, urine, food, water, and culture inoculated with at least one of blood, sputum, stool, tissue, media, semen, hair, urine, food, and water.

25

15

10. The method of claim 1 wherein the nucleic acid of interest is derived from a source selected from the group consisting of a bacterium, a virus, a human, an animal, or a plant.

30

20

11. The method according to claim 1 wherein at least one primer of any one of said sets is in molar excess as compared to any other primer in such set.

35

12. The method according to claim 11 wherein a primer that is in molar excess in said amplification reaction is labeled.

40

25

13. A method for the amplification, multiplex assaying, and detection of target nucleic acids of interest using a bioelectronic microchip comprising the steps of:

45

- a. introducing at least one of said target nucleic acids of interest onto a bioelectronic microchip having a plurality of electronically addressable capture sites;

30

50

55

5

10

5

15

20

10

25

15

30

20

35

40

25

45

30

50

55

- b. addressing said target nucleic acids of interest to at least one of said capture sites;
- c. amplifying said target nucleic acids to form amplicons of said target nucleic acids wherein said target nucleic acids are amplified using an amplification reaction wherein a multiplicity of sets of amplification primers are used that include unequal ratios of upstream and downstream primers;
- d. capturing said amplicons of said target nucleic acids onto specified capture sites of said microchip by a capture probe that has specificity for said amplicons; and
- e. detecting the presence of said captured amplicons.

14. The method of claim 13 wherein amplification of the target nucleic acid is by strand displacement amplification.

15. The method of claim 13 wherein amplification of the target nucleic acid is by at least one of the methods selected from the group consisting of strand displacement amplification, allele-specific strand displacement amplification, nucleic acid sequence-based amplification, and ligation-based strand displacement amplification.

16. The method of claim 13 wherein said addressing of said target nucleic acids in (b) is carried out by passive hybridization.

17. The method of claim 13 wherein said addressing of said target nucleic acids in (b) is carried out by electronic biasing of said capture sites.

18. The method of claim 13 wherein said capturing of amplicons in (d) is carried out by passive hybridization.

19. The method of claim 13 wherein said capturing of amplicons in (d) is carried out by electronic biasing of said capture sites.

5

20. The method of claim 13 wherein said amplification is carried out in part using noncleavable primers and/or non-extending primers.

10

21. The method of claim 13 wherein said amplification, multiplex assaying and detecting are carried out either consecutively or simultaneously in relation to one another.

5

15

22. The method of claim 13 wherein detection of amplicons is by at least one of fluorescence, chemiluminescence, and electrochemiluminescence.

20

23. The method of claim 13 wherein target nucleic acids of interest are derived from a source selected from the group consisting of blood, sputum, stool, tissue, media, semen, hair, urine, food, water, and culture inoculated with at least one of blood, sputum, stool, tissue, media, semen, hair, urine, food, and water.

25

24. The method of claim 13 wherein the nucleic acid of interest is derived from a source selected from the group consisting of a bacterium, a virus, a human, an animal, or a plant.

30

25. The method according to claim 13 wherein at least one primer of any one of said sets is in molar excess as compared to any other primer in such set.

20

35

26. The method according to claim 25 wherein a primer that is in molar excess in said amplification reaction is labeled.

40

27. A kit for carrying out asymmetric SDA reactions for use on a bioelectronic microchip comprising:

25

- a. one or more oligonucleotides specific for Factor V, Hemochromatosis, or a bacterium, which oligonucleotides comprise amplification primers, bumper primers, capture probes, and/or signal probes selected from the group consisting of Seq. Id. Nos. 1-62.

45

30

50

55

5

28. A kit according to claim 27 wherein said signal probes are labeled with a detectable label.

10

15

20

25

30

35

40

45

50

55

1/34




FIGURE 1A





FIGURE 1B


2/34

FIGURE 2A



3/34

FIGURE 2B



4/34

FIGURE 2D






FIGURE 2C






5/34




6/34

FIGURE 3B




**FIGURE 3C**



Salmonella Target DNA Input/Universal Reporter


8/34

FIGURE 4A




9/34

FIGURE 4B




10/34


FIGURE 5A



11/34


FIGURE 5B








13/34




14/34

FIGURE 6C




15/34


**FIGURE 7**

1005 + 1006 = monobiotinylated probes  
1877 + 1878 = quadruple-biotinylated probes  
1029 + 1030 = non-cleavable control primers

16/34




17/34

**FIGURE 9**

1005 + 1006 = Factor V SDA primers  
1029 + 1030 = Non-cleavable control primers


18/34

FIGURE 10A




19/34

FIGURE 10B



20/34


FIGURE 11






21/34

**FIGURE 12**




22/34




23/34

**FIGURE 14**



24/34

**FIGURE 15**




25/34

FIGURE 17



Control - No template +  
all reporter oligos


FIGURE 16




Experimental Layout

**FIGURE 18**

**FIGURE 19**




**FIGURE 20**

28/34

**FIGURE 21**

Factor V	Chlamydia	Hemochrom	ALL
-	+	-	+
+	-	+	-




Control Solution SDA reactions



29/34

FIGURE 22






Fig. 23 A

31/34





Fig. 23 B

32/34



(Product XII can re-enter pathway and be further amplified in a manner similar to product III, following step 6)

Fig. 23 C

Fig. 23 D Ligation-dependent detection of the *Salmonella* spaQ gene.

## Ligation Probes LP1 and LP2:

spaQ template 5'-nnnnncaacatgacatcattacgagacgggatagttaaatggatgatttagtgmnnn-3'  
 LP1 3'-aattccgcatgagctgggtaagtgttactgtagtaatgctctgc\*-5'  
 LP2 3'-cctatcaattacactactaataacacgattatccctagagatgcatgtggctc ttcagacctgccttagc-5'


## Amplification primer sequences S1 and S2:

S1 5'-accgcatcgaaatgcattgctctcgggtaaggctactcgacc  
 LP1 3'-aattccgcatgagctgggtaagtgttactgtagtaatgctctgc\*-5'  
 S2 3'-cctatcaattacactactaataacacgattatccctagagatgcatgtggctccttcagacctgccttagc-5'

\* spaQ (Seq. ID #41)  
 \* LP1 (Seq. ID #37); \* = phosphate;  
 5' phosphate on LP1 provides ligatable end  
 3' phosphate on LP1 blocks polymerase extension  
 \* LP2 (Seq. ID #38)  
 \* S1 (Seq. ID #39)  
 \* S2 (Seq. ID #40)

34/34

Fig. 24



Electronic hybridization using microelectrode array of XL-SDA reactions with Salmonella genomic DNA at 1000

## SEQUENCE LISTING

5 <110> Nerenberg, Michael I.  
Westin, Lorelei P.  
Edman, Carl F.  
Carrino, John

10 <120> MULTIPLEX AMPLIFICATION AND SEPARATION OF NUCLEIC ACID  
SEQUENCES ON A BIOELECTRONIC MICROCHIP USING  
ASYMMETRIC  
STRUCTURES

15 <130> 241/109

<140> TO BE ASSIGNED  
<141> HERewith

20 <160> 62

<170> FastSEQ for Windows Version 3.0

25 <210> 1  
<211> 20  
<212> DNA  
<213> conserved 16S bacterial sequence

30 <400> 1

caaatgaatt gacgggggcc

20

<210> 2  
<211> 15  
<212> DNA  
5 <213> conserved 16S bacterial sequence

<400> 2

10 aagggttgcg ctcgt

15

<210> 3  
<211> 40  
15 <212> DNA  
<213> conserved 16S bacterial sequence

<400> 3

20 accgcatoga atgcatgtcc tcgggtgcat gtggtttaat

40

<210> 4  
25 <211> 41  
<212> DNA  
<213> conserved 16S bacterial sequence

<400> 4

30 acgattcagc tccagacttc tcgggtaaca ttccacaaca c

41



5 <210> 5  
<211> 20  
<212> DNA  
<213> human

<400> 5

10 actacagtga cgtggacatc 20

15 <210> 6  
<211> 20  
<212> DNA  
<213> human

<400> 6

20 tgttatcaca ctggtgctaa 20

25 <210> 7  
<211> 42  
<212> DNA  
<213> human

30 <400> 7

accgcacga atgcatgtcc tcgggtctct gggctaataag ga 42

<210> 8  
<211> 42  
5 <212> DNA  
<213> human

<400> 8

10 acgattcagc tccagacttc tcgggtaata cctgtattcc tc

42

<210> 9  
15 <211> 42  
<212> DNA  
<213> human

<400> 9

20

acgattcagc tccagacttc tcgggtaata cctgtattcc tt

42

25 <210> 10  
<211> 18  
<212> DNA  
<213> human

30 <400> 10

ctgtattcct cgcctgtc

18

	<210> 11	
	<211> 18	
5	<212> DNA	
	<213> E. coli	
	<400> 11	
10	ctcatctctg aaaacttc	18
	<210> 12	
15	<211> 18	
	<212> DNA	
	<213> Shigella dysenteriae	
	<400> 12	
20	cgtaactcta caaggttc	18
	<210> 13	
25	<211> 18	
	<212> DNA	
	<213> Salmonella typhimurium	
30	<400> 13	
	tccatctctg gattcttc	18

<210> 14  
<211> 18  
5 <212> DNA  
<213> Camphylobacter Jejuni  
  
<400> 14  
10 catatctcta taaggttc 18  
  
<210> 15  
15 <211> 27  
<212> DNA  
<213> conserved 16S bacterial sequence  
  
<400> 15  
20 ggatgtcaag accaggtaag gttcttc 27  
  
<210> 16  
25 <211> 50  
<212> DNA  
<213> human T-cell leukemia virus-1  
  
<400> 16  
30 aattctaata cgactcacta tagggagagg tgatctgatg tctggacagg 50

5           <210> 17  
            <211> 23  
            <212> DNA  
            <213> human T-cell leukemia virus-1

            <400> 17

10       acttcccagg gtttggacag agt

23

15           <210> 18  
            <211> 30  
            <212> DNA  
            <213> human T-cell leukemia virus-1

            <400> 18

20       ttcttttcgg ataccagtc tacgtgttg

30

25           <210> 19  
            <211> 23  
            <212> DNA  
            <213> human T-cell leukemia virus-1

30           <400> 19

acttcccagg gtttggacag agt

23

5           <210> 20  
            <211> 42  
            <212> DNA  
            <213> human

            <400> 20

10           accgcatcga atgcatgtcc tcgggtctct gggctaataag ga

42

15           <210> 21  
            <211> 42  
            <212> DNA  
            <213> human

            <400> 21

20

            acgattcagc tccagacttc tcgggtcaga atttctgaaa gg

42

25           <210> 22  
            <211> 20  
            <212> DNA  
            <213> human

30

            <400> 22

            actacagtga cgtggacatc

20

5 <210> 23  
<211> 20  
<212> DNA  
<213> human

<400> 23

10 tggtatcaca ctggtgctaa

20

15 <210> 24  
<211> 18  
<212> DNA  
<213> human

<400> 24

20 ctgtattcct egcctgtc

18

25 <210> 25  
<211> 42  
<212> DNA  
<213> Chlamydia trachomatis

30 <400> 25

cacgtagtca atgcatgtcc tcggtacaa catcaacacc tg

42

5           <210> 26  
            <211> 42  
            <212> DNA  
            <213> Chlamydia trachomatis

            <400> 26

10       acgattcagc tccagacttc tcgggtgaga ctgttaaaga ta

42

15           <210> 27  
            <211> 17  
            <212> DNA  
            <213> Chlamydia trachomatis

            <400> 27

20       cagcaaataa tccttgg

17

25           <210> 28  
            <211> 19  
            <212> DNA  
            <213> Chlamydia trachomatis

30           <400> 28

cattggttga tggattatt

19



5           <210> 29  
            <211> 15  
            <212> DNA  
            <213> Chlamydia trachomatis

            <400> 29

10       gtcgcagcca aaatg

15

15           <210> 30  
            <211> 16  
            <212> DNA  
            <213> Chlamydia trachomatis

            <400> 30

20

ttccatcaga agctgt

16

25           <210> 31  
            <211> 42  
            <212> DNA  
            <213> human

30

            <400> 31

cacgtagtca atgcatgtcc tcgggtataa ccttggtgt ac

42

5           <210> 32  
            <211> 42  
            <212> DNA  
            <213> human

            <400> 32

10       acgattcagc tccagacttc tgggtgctc tcacagtca ca

42

15           <210> 33  
            <211> 19  
            <212> DNA  
            <213> human

            <400> 33

20

      tgaaggataa gcagccaat

19

25           <210> 34  
            <211> 19  
            <212> DNA  
            <213> human

30

            <400> 34

      ctctctcaaa cccccaata

19

5       <210> 35  
          <211> 18  
          <212> DNA  
          <213> human

          <400> 35

10       agatatacgt gccaggtg

18

15       <210> 36  
          <211> 18  
          <212> DNA  
          <213> human

          <400> 36

20       ctgatccagg cctgggtg

18

25       <210> 37  
          <211> 45  
          <212> DNA  
          <213> Salmonella

30       <400> 37

aattccgat gagctgggta atgtgtact gtagtaatgc tctgc

45

5 <210> 38  
<211> 70  
<212> DNA  
<213> Salmonella

<400> 38

10 cctatcaatt tacctactaa atcacgatta tcccctagag tcatgtgggc tcttcagacc 60  
tcgccttagc 70

15 <210> 39  
<211> 40  
<212> DNA  
<213> synthetic

20 <400> 39

accgcatcga atgcatgtct cgggtaaggc gtactcgacc 40

25 <210> 40  
<211> 40  
<212> DNA  
<213> synthetic

30 <400> 40

cgattccgct ccagacttct cgggtgtact gagatcccct 40

5           <210> 41  
            <211> 48  
            <212> DNA  
            <213> synthetic  
  
            <400> 41  
10           caacatgaca tcattacgag acgggatagt taaatggatg atttagtg           48  
  
            <210> 42  
15           <211> 42  
            <212> DNA  
            <213> human  
  
            <400> 42  
20           accgcatcga atgcatgtcc tccggtctct gggctaataag ga           42  
  
            <210> 43  
25           <211> 42  
            <212> DNA  
            <213> human  
  
            <400> 43  
30           acgattcaga tccagacttc tccggtcaga atttctgaaa gg           42

5           <210> 44  
            <211> 21  
            <212> DNA  
            <213> human

            <400> 44

10       acttctaact tgtaagagca g

21

15           <210> 45  
            <211> 41  
            <212> DNA  
            <213> synthetic

            <400> 45

20       gagggcggtt taataatcta cgggtggtcga gtacgcctta a

41

25           <210> 46  
            <211> 63  
            <212> DNA  
            <213> synthetic

30           <400> 46

cgattccgct ccagacttct cgggtgtact gagatcccct tgcagaggg atagatccag  
agg

60

63

5 <210> 47  
<211> 43  
<212> DNA  
<213> synthetic

<400> 47

10 gatggagttc agtggtaata caatgtggtc gactacgcct taa

43

15 <210> 48  
<211> 61  
<212> DNA  
<213> synthetic

<400> 48

20 cgattccgct ccagacttct cgggtgtact gagatcccct ggtttcatca tatctggcgt  
t

60

61

25 <210> 49  
<211> 39  
<212> DNA  
<213> synthetic

30

<400> 49

gacgtgctc actagatgtc taggtcgagt acgccttaa

39

	<210> 50	
	<211> 64	
5	<212> DNA	
	<213> synthetic	
	<400> 50	
10	cgattccgct ccagacttct cgggtgtact gagatcccct ggttataagt gcttgatact	60
	ccag	64
	<210> 51	
	<211> 41	
15	<212> DNA	
	<213> synthetic	
	<400> 51	
20	gatgatgtca tgttgcaatg tcctggtcga gtacgcccta a	41
	<210> 52	
25	<211> 61	
	<212> DNA	
	<213> synthetic	
	<400> 52	
30	cgattccgct ccagacttct cgggtgtact gagatcccct catttaacta tcccgtctcg	60
	t	61



5           <210> 53  
            <211> 44  
            <212> DNA  
            <213> synthetic

            <400> 53

10       gagtaattac cgtcttcac ttttttggc cgagtacgcc ttaa

44

15           <210> 54  
            <211> 65  
            <212> DNA  
            <213> synthetic

            <400> 54

20

cgattccgct ccagacttct cgggtgtact gagatccctt ggcttcacaa agaataacat  
ctatc

60

65

25           <210> 55  
            <211> 40  
            <212> DNA  
            <213> synthetic

            <400> 55

30

gatttacgga ctggtctcc ctgggtcgag tacgccttaa

40

<210> 56  
<211> 61  
<212> DNA  
5 <213> synthetic

<400> 56

cgattccgct ccagacttct cgggtgtact gagatcccct tcagaagccg tgaagagaat 60  
10 g 61

<210> 57  
15 <211> 39  
<212> DNA  
<213> synthetic

<400> 57

20 gacccaaaacc atcctgaacc atggtcgagt acgccttaa 39

25 <210> 58  
<211> 67  
<212> DNA  
<213> synthetic

30 <400> 58

cgattccgct ccagacttct cgggtgtact gagatcccct ttctagtttt tgatttttag 60  
tattata 67

5 <210> 59  
<211> 42  
<212> DNA  
<213> synthetic

<400> 59

10 gagtagaggt atgtgatgag ccaatggcgc agtacgcctt aa 42

15 <210> 60  
<211> 61  
<212> DNA  
<213> synthetic

<400> 60

20 cgattccgct ccagacttct cgggtgtact gagatcccct ctttgctaa actcggtttt 60  
c 61

25 <210> 61  
<211> 39  
<212> DNA  
<213> synthetic

30 <400> 61

gattagctga gcttaccgcc gtggcgcagt acgccitaa 39

&lt;210&gt; 62

&lt;211&gt; 60

5 &lt;212&gt; DNA

&lt;213&gt; synthetic

&lt;400&gt; 62

10 cgattccgct ccagacttct cgggigtact gagatcccct ccgtagcaag ttgcgtgaag 60

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/09742

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(7) : C12Q 1/68 US CL : 435/6 According to International Patent Classification (IPC) or to both national classification and IPC														
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/6  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAPLUS BIOSIS MEDLINE SCISEARCH USPATFUL														
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>														
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
Y	WO 96/01836 A1 (HELLER et al) 25 January 1996, page 9 lines 13-16, especially pages 13-15, page 16 lines 27-29.	1-28												
Y	SPARGO, C.A. Detection of M. tuberculosis DNA using Thermophilic Strand Displacement Amplification, Molecular and Cellular Probes. 1996, Vol. 10, pages 247-256, see entire document.	1-28												
Y	US 5,593,867 A (WALKER et al) 14 January 1997, column 6, lines 45-49.	12, 26												
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.														
<table border="0"><tr><td>* Special categories of cited documents:</td><td>* "T" later document published after the international filing date or priority date and not in conflict with the application but aimed to understand the principle or theory underlying the invention</td></tr><tr><td>* "A" document defining the general state of the art which is not considered to be of particular relevance</td><td>* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td></tr><tr><td>* "B" earlier document published on or after the international filing date</td><td>* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combinations being obvious to a person skilled in the art</td></tr><tr><td>* "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td><td>* "A" document member of the same patent family</td></tr><tr><td>* "O" document referring to an oral disclosure, use, exhibition or other means</td><td></td></tr><tr><td>* "P" document published prior to the international filing date but later than the priority date claimed</td><td></td></tr></table>			* Special categories of cited documents:	* "T" later document published after the international filing date or priority date and not in conflict with the application but aimed to understand the principle or theory underlying the invention	* "A" document defining the general state of the art which is not considered to be of particular relevance	* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	* "B" earlier document published on or after the international filing date	* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combinations being obvious to a person skilled in the art	* "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* "A" document member of the same patent family	* "O" document referring to an oral disclosure, use, exhibition or other means		* "P" document published prior to the international filing date but later than the priority date claimed	
* Special categories of cited documents:	* "T" later document published after the international filing date or priority date and not in conflict with the application but aimed to understand the principle or theory underlying the invention													
* "A" document defining the general state of the art which is not considered to be of particular relevance	* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone													
* "B" earlier document published on or after the international filing date	* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combinations being obvious to a person skilled in the art													
* "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* "A" document member of the same patent family													
* "O" document referring to an oral disclosure, use, exhibition or other means														
* "P" document published prior to the international filing date but later than the priority date claimed														
Date of the actual completion of the international search 17 JULY 2000		Date of mailing of the international search report 15 AUG 2000												
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer <i>Jeffrey S. Landgren</i> Jeffrey S. Landgren Telephone No. (703) 308-0196												